

THE CUNTZ SEMIGROUP OF CONTINUOUS FIELDS

RAMON ANTOINE, JOAN BOSCH, AND FRANCESC PERERA

ABSTRACT. In this paper we describe the Cuntz semigroup of continuous fields of C^* -algebras over one dimensional spaces whose fibers have stable rank one and trivial K_1 for each closed, two-sided ideal. This is done in terms of the semigroup of global sections on a certain topological space built out of the Cuntz semigroups of the fibers of the continuous field. When the fibers have furthermore real rank zero, and taking into account the action of the space, our description yields that the Cuntz semigroup is a classifying invariant if and only if so is the sheaf induced by the Murray-von Neumann semigroup.

INTRODUCTION

The Cuntz semigroup has become a popular object in recent years, mainly due to its connection with the classification program of unital, simple, separable and nuclear C^* -algebras by means of the Elliott invariant. It can be thought of as a functorial invariant $\text{Cu}(\cdot)$ from the category of C^* -algebras to a category of semigroups, termed Cu , that has certain continuity properties. The Elliott invariant is also functorial and consists of ordered topological K-Theory, the trace simplex and the pairing between K-Theory and traces; it is customarily denoted by $\text{Ell}(\cdot)$. The Elliott conjecture asserts that an isomorphism between the invariants $\text{Ell}(A)$ and $\text{Ell}(B)$ of C^* -algebras A and B may be lifted to a $*$ -isomorphism between A and B . Although the conjecture fails in general (see [26], [29]), it has been verified for large classes of C^* -algebras, all of which happen to absorb the Jiang-Su algebra \mathcal{Z} tensorially. For these algebras, the Cuntz semigroup can be recovered functorially from the Elliott invariant (see [7]). More recently, it was shown further in [2] (see also [28]) that the Elliott invariant can be recovered from the Cuntz semigroup after tensoring with $C(\mathbb{T})$, and thus $\text{Ell}(\cdot)$ and $\text{Cu}(C(\mathbb{T}), \cdot)$ define equivalent functors for these algebras.

In the non-simple case, the Cuntz semigroup has already been used successfully to classify certain classes of C^* -algebras, such as AI algebras ([9]), inductive limits of one dimensional non commutative CW-complexes with trivial K_1 ([23]) or inductive limits of certain continuous-trace C^* -algebras ([10]), among others. Another class of (non-simple) algebras for which classification results have been obtained are continuous fields over $[0, 1]$ of either Kirchberg algebras (with certain torsion freeness assumptions on their K-Theory) or AF algebras ([13, 14]). In this situation, the classifying invariant consists of the sheaf of groups naturally induced by K-Theory. In the stably finite case, it is natural to ask whether the Cuntz semigroup of the continuous field captures, on its own, all the information of the K-Theory sheaf. This is one of the main objectives pursued in this article, and we are able to settle the question positively for a wide class of continuous fields.

In order to achieve our aim, we need techniques that allow us to compute the Cuntz semigroup of continuous fields. In the case of algebras of the form $C_0(X, A)$, for a locally

compact Hausdorff space X , this was carried out in [28] whenever A is a simple, unital, non-type I ASH-algebra with slow dimension growth. For a not necessarily simple algebra A of stable rank one with vanishing K_1 for each closed, two-sided ideal, and for a one dimensional locally compact Hausdorff space, one of the main results in [3] proves that there is an isomorphism between $\text{Cu}(C_0(X, A))$ and $\text{Lsc}(X, \text{Cu}(A))$. The latter is a semigroup of $\text{Cu}(A)$ -valued lower semicontinuous functions (see below for the precise definitions), and the isomorphism is given by point evaluation of (representatives of) Cuntz classes. The more general situation of $C(X)$ -algebras was also a theme developed in [3], where spaces of dimension at most one and $C(X)$ -algebras whose fibers have stable rank one and vanishing K_1 for each closed, two-sided ideal were considered. In this paper, we shall refer to this class as *$C(X)$ -algebras without K_1 obstructions*. For such spaces, the Cuntz semigroup of a $C(X)$ -algebra A with no K_1 obstructions embeds into the product $\prod_{x \in X} \text{Cu}(A_x)$. (This was shown in [3] for $X = [0, 1]$, and we prove the one dimensional case here, based on the pullback construction carried out also in [3].)

The remaining problem of identifying the image of $\text{Cu}(A)$ in $\prod_{x \in X} \text{Cu}(A_x)$ for $C(X)$ -algebras A in the said class leads to the analysis of the natural map $F_{\text{Cu}(A)} := \sqcup_{x \in X} \text{Cu}(A_x) \rightarrow X$ and its sections. This is motivated by the fact that the Cu functor induces a presheaf Cu_A on X that assigns, to each closed set U of X , the semigroup $\text{Cu}(A(U))$. Hence we may expect to relate the Cuntz semigroup with the semigroup of continuous sections of an étale bundle. In the case of the presheaf defined by K-Theory, and for some continuous fields over $[0, 1]$, this was considered in [13]. We show that, for a one dimensional space X and a $C(X)$ -algebra with no K_1 obstructions, the presheaf Cu_A is in fact a sheaf. In order to recover Cu_A from the sheaf of continuous sections of the map $F_{\text{Cu}(A)} \rightarrow X$, we need to break away from the standard approach (of, e.g. [30]) and consider a topological structure on $F_{\text{Cu}(A)}$ that takes into account continuity properties of the objects in the category Cu . Thus we develop a more abstract analysis of Cu -valued sheaves that follows, in part, the spirit of [3]. This culminates in Theorem 3.12, which allows to recover the Cuntz semigroup of a $C(X)$ -algebra A with no K_1 obstructions over a one dimensional space X as the semigroup of global sections on $F_{\text{Cu}(A)}$.

To conclude the paper, we apply the previous result in a crucial way to prove that, for one dimensional spaces and $C(X)$ -algebras with no K_1 obstructions whose fibers have real rank zero, the Cuntz semigroup and the K -theoretical sheaf defined by the Murray-von Neumann semigroup carry the same information. (This sheaf is defined, for a $C(X)$ -algebra A , as $\mathbb{V}_A(U) = V(A(U))$ whenever U is a closed subset of X .) A key ingredient here is that the natural module structure of a $C(X)$ -algebra A equips $\text{Cu}(A)$ with an enriched structure via an action of $\text{Cu}(C(X))$. Thus, more precisely, we show that if A and B are two such $C(X)$ -algebras, then there is an action preserving semigroup isomorphism $\text{Cu}(A) \cong \text{Cu}(B)$ if, and only if, the sheaves $\mathbb{V}_A(-)$ and $\mathbb{V}_B(-)$ are isomorphic (Theorem 4.9).

1. PRELIMINARIES

1.1. Cuntz Semigroup. Let A be a C^* -algebra, and let $a, b \in A_+$. We say that a is *Cuntz subequivalent* to b , in symbols $a \preceq b$, provided there is a sequence (x_n) in A such that $x_n b x_n^*$ converges to a in norm. We say that a and b are *Cuntz equivalent* if $a \preceq b$ and $b \preceq a$, and in this case we write $a \sim b$.

The Cuntz semigroup is defined as the quotient set $\text{Cu}(A) = (A \otimes \mathcal{K})_+ / \sim$, and its elements are denoted by $[a]$, for $a \in (A \otimes \mathcal{K})_+$. This set becomes an ordered semigroup, with order induced by Cuntz subequivalence and addition given by $[a] + [b] = [\Theta \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}]$, where $\Theta: M_2(A \otimes \mathcal{K}) \rightarrow A \otimes \mathcal{K}$ is any isomorphism. The following summarizes some technical properties of Cuntz subequivalence that will be used in the sequel.

Proposition 1.1. ([25], [16]) *Let A be a C^* -algebra, and $a, b \in A_+$. The following are equivalent:*

- (i) $a \preceq b$.
- (ii) For all $\epsilon > 0$, $(a - \epsilon)_+ \preceq b$.
- (iii) For all $\epsilon > 0$, there exists $\delta > 0$ such that $(a - \epsilon)_+ \preceq (b - \delta)_+$.
Furthermore, if A is stable, these conditions are equivalent to
- (iv) For every $\epsilon > 0$ there is a unitary $u \in U(\tilde{A})$ such that $u(a - \epsilon)_+ u^* \in \text{Her}(b)$.

The structure of the Cuntz semigroup is richer than just being an ordered semigroup, as it belongs to a category with certain continuity properties. Recall that, in an ordered semigroup S , an element s is said to be *compactly contained* in t , denoted $s \ll t$, if whenever $t \leq \sup_n z_n$ for some increasing sequence (z_n) with supremum in S , there exists m such that $s \leq z_m$. An element s is said to be *compact* if $s \ll s$. A sequence (s_n) such that $s_n \ll s_{n+1}$ is termed *rapidly increasing*. The following theorem summarizes some structural properties of the Cuntz semigroup.

Theorem 1.2. ([8]) *Let A be a C^* -algebra. Then:*

- (i) Every increasing sequence in $\text{Cu}(A)$ has a supremum in $\text{Cu}(A)$.
- (ii) Every element in $\text{Cu}(A)$ is the supremum of a rapidly increasing sequence.
- (iii) The operation of taking suprema and \ll are compatible with addition.

This allows one to define a category Cu whose objects are ordered semigroups of positive elements satisfying conditions (i)-(iii) above. (Morphisms in this category are those semigroup maps that preserve all the structure.) We say that a semigroup S in the category Cu is *countably based* if there exists a countable subset X that is dense in S , meaning that every element of S is the supremum of a rapidly increasing sequence of elements in X . It was observed in [3] (see also [22]) that $\text{Cu}(A)$ is countably based for any separable C^* -algebra A .

As shown in [8], the category Cu is closed under countable inductive limits (in fact, it was also shown that Cu defines a sequentially continuous functor from the category of C^* -algebras to Cu). A useful description of the inductive limit is available below.

Proposition 1.3. ([8], cf. [5]) *Let $(S_i, \alpha_{i,j})_{i,j \in \mathbb{N}}$ be an inductive system in the category Cu . Then $(S, \alpha_{i,\infty})$ is the inductive limit of this system if*

- (i) The set $\bigcup_i \alpha_{i,\infty}(S_i)$ is dense in S .
- (ii) For any $x, y \in S_i$ such that $\alpha_{i,\infty}(x) \leq \alpha_{i,\infty}(y)$ and $x' \ll x$ there is j such that $\alpha_{i,j}(x') \leq \alpha_{i,j}(y)$.

If S is a semigroup in Cu , an *order-ideal* I of S is a subsemigroup which is order-hereditary (that is, $x \in I$ whenever $x \leq y$ and $y \in I$) and that further contains all suprema of increasing sequences in I . For example, if A is a C^* -algebra and I is a closed, two-sided ideal, then $\text{Cu}(I)$ is naturally an order-ideal of $\text{Cu}(A)$. Given an order-ideal I of S as before, define a congruence relation on S by $s \sim t$ if $s \leq t + z$ and $t \leq s + w$ for some $z, w \in I$, and

put $S/I = S/\sim$, which is an ordered semigroup with addition $[s] + [t] = [s + t]$ and order given by $[s] \leq [t]$ if $s \leq t + z$ for some $z \in I$. It is not hard to verify that $S/I \in \text{Cu}$. For a C^* -algebra A and a closed ideal I , it was proved in [11] that $\text{Cu}(A/I) \cong \text{Cu}(A)/\text{Cu}(I)$ (where the isomorphism is induced by the natural quotient map).

1.2. $C(X)$ -algebras. Let X be a compact Hausdorff space. A $C(X)$ -algebra is a C^* -algebra A together with a unital $*$ -homomorphism $\theta: C(X) \rightarrow Z(\mathcal{M}(A))$, where $\mathcal{M}(A)$ is the multiplier algebra of A . The map θ is usually referred to as the *structure map*. We write fa instead of $\theta(f)a$ where $f \in C(X)$ and $a \in A$.

If $Y \subseteq X$ is a closed set, let $A(Y) = A/C_0(X \setminus Y)A$, which also becomes a $C(X)$ -algebra. The quotient map is denoted by $\pi_Y: A \rightarrow A(Y)$, and if Z is a closed subset of Y we have a natural restriction map $\pi_Z^Y: A(Y) \rightarrow A(Z)$. Notice that $\pi_Z = \pi_Z^Y \circ \pi_Y$. If Y reduces to a point x , we write A_x instead of $A(\{x\})$ and we denote by π_x the quotient map. The C^* -algebra A_x is called the *fiber* of A at x and the image of $\pi_x(a) \in A_x$ will be denoted by $a(x)$.

Given a $C(X)$ -algebra A and $a \in A$, the map $x \mapsto \|a(x)\|$ is upper semicontinuous (see [6]). If this map is actually continuous for every $a \in A$, then we say that A is a *continuous field* (or also a C^* -bundle, see [17, 6]). For a continuous field A , a useful criterion to determine when an element $(a_x) \in \prod_{x \in X} A_x$ comes from an element of A is the following: given $\epsilon > 0$ and $x \in X$, if there is $b \in A$ and a neighborhood V of x such that $\|b(y) - a_y\| < \epsilon$ for $y \in V$, then there is $a \in A$ such that $a(x) = a_x$ for all x (see [15, Definition 10.3.1]).

It was proved in [3, Lemma 1.5] that, if A is a $C(X)$ -algebra, then this is also the case for $A \otimes \mathcal{K}$ and, in fact, for any closed set Y of X , there is a $*$ -isomorphism

$$\varphi_Y: (A \otimes \mathcal{K})(Y) \rightarrow A(Y) \otimes \mathcal{K}$$

such that $\varphi_Y \circ \pi_Y' = \pi_Y \otimes 1_{\mathcal{K}}$, where $\pi_Y: A \rightarrow A(Y)$ and $\pi_Y': A \otimes \mathcal{K} \rightarrow (A \otimes \mathcal{K})(Y)$. This yields, in particular, that $(A \otimes \mathcal{K})(x) \cong A_x \otimes \mathcal{K}$ for any $x \in X$, with $(a \otimes k)(x) \mapsto a(x) \otimes k$.

Using this observation, the map induced at the level of Cuntz semigroups $\text{Cu}(A) \rightarrow \text{Cu}(A_x)$ can be viewed as $[a] \mapsto [\pi_x(a)]$. Similarly, if Y is closed in X , the map π_Y induces $\text{Cu}(A) \rightarrow \text{Cu}(A(Y))$, that can be thought of as $[a] \mapsto [\pi_Y(a)]$. Thus, when computing the Cuntz semigroup of a $C(X)$ -algebra A , we may and will assume that A , A_x and $A(Y)$ are stable.

2. SHEAVES OF SEMIGROUPS AND CONTINUOUS SECTIONS

Our aim in this section is to relate the Cuntz semigroup of a $C(X)$ -algebra A with the semigroup of continuous sections of a certain topological space, which is built out of the information on the fibers. We will first define what is meant by a presheaf of semigroups on a topological space, along the lines of [30], with some modifications.

Let X be a topological space. As a blanket assumption, we shall assume that X is always compact, Hausdorff and second countable, therefore metrizable. Denote by \mathcal{V}_X the category of all closed subsets of X with non-empty interior, with the morphisms given by inclusion.

A *presheaf* over X is a contravariant functor

$$S: \mathcal{V}_X \rightarrow \mathcal{C}$$

where \mathcal{C} is a (not necessarily full) subcategory of the category of sets such that \mathcal{C} is closed under sequential inductive limits. In practice, \mathcal{C} will be the category Sg of semigroups

(with semigroup maps) or the category Cu (with morphisms in Cu as defined before). Thus, a presheaf over X consists of an assignment, for each $V \in \mathcal{V}_X$ of an object $\mathcal{S}(V)$ in \mathcal{C} and a collection of maps (referred to as *restriction homomorphisms*) $\pi_V^{V'} : \mathcal{S}(V') \rightarrow \mathcal{S}(V)$ whenever $V \subseteq V'$ in \mathcal{V}_X . We of course require that these maps satisfy $\pi_V^V = \text{id}_V$ and $\pi_W^U = \pi_W^V \pi_V^U$ if $W \subseteq V \subseteq U$.

Let $V, V' \in \mathcal{V}_X$ be such that $V \cap V' \in \mathcal{V}_X$. A presheaf is called a *sheaf* if the map

$$\pi_V^{V \cup V'} \times \pi_{V'}^{V \cup V'} : \mathcal{S}(V \cup V') \rightarrow \{(f, g) \in \mathcal{S}(V) \times \mathcal{S}(V') \mid \pi_{V \cap V'}^V(f) = \pi_{V \cap V'}^{V'}(g)\},$$

is bijective.

A presheaf (respectively a sheaf) is *continuous* if for any decreasing sequence of closed subsets $(V_i)_{i=1}^\infty$ whose intersection $\bigcap_{i=1}^\infty V_i = V$ belongs to \mathcal{V}_X , the limit $\lim \mathcal{S}(V_i)$ is isomorphic to $\mathcal{S}(V)$.

Consider a presheaf \mathcal{S} over X . For any $x \in X$, define the *fiber* (or also *stalk*) of \mathcal{S} at x as

$$S_x := \lim_{x \in V} \mathcal{S}(V),$$

with respect to the restriction maps.

We shall be exclusively concerned with continuous presheaves (or sheaves) \mathcal{S} with target values in the category Sg of semigroups, in which case we will say that \mathcal{S} is a (pre)sheaf of semigroups. As a general notation, we will use S to denote the semigroup $\mathcal{S}(X)$. We will also denote $\pi_x : S \rightarrow S_x$ the natural map from S to the fiber S_x , as well as $\pi_U : S \rightarrow S(U)$ rather than π_U^X .

Our main motivation for considering presheaves of semigroups stems from the study of $C(X)$ -algebras. Indeed, as it is easy to verify, given a $C(X)$ -algebra A , the assignments

$$\text{Cu}_A : \begin{array}{ccc} \mathcal{V}_X & \rightarrow & \text{Cu} \\ U & \mapsto & \text{Cu}(A(U)) \end{array} \quad \text{and} \quad \mathbb{V}_A : \begin{array}{ccc} \mathcal{V}_X & \rightarrow & \text{Sg} \\ U & \mapsto & V(A(U)) \end{array}$$

define continuous presheaves of semigroups. If $U \subseteq V$, the restriction maps $\pi_V^U : A(U) \rightarrow A(V)$ and the limit maps $\pi_x : A \rightarrow A_x$ define, by functoriality, semigroup maps $\text{Cu}(\pi_V^U)$ and $\text{Cu}(\pi_x)$ in the case of the Cuntz semigroup, and likewise in the case for the semigroup of projections. For ease of notation, and unless confusion may arise, we shall still denote these maps by π_V^U and π_x .

We will say that a (pre)sheaf is *surjective* provided all the restriction maps are surjective. This is clearly the case for the presheaf Cu_A for a general $C(X)$ -algebra A , and also for \mathbb{V}_A if A has real rank zero (which is a rather restrictive hypothesis, see e.g. [18] and [19]). As we shall see in the sequel, Cu_A and \mathbb{V}_A determine each other under milder assumptions.

Most of the discussion in this and the subsequent section will consider surjective (pre)sheaves of semigroups $\mathcal{S} : \mathcal{V}_X \rightarrow \text{Cu}$, and we will need to develop a somewhat abstract approach on how to recover the information of the sheaf from the sheaf of sections of a bundle $F_S \rightarrow X$, where F_S stands for the disjoint union of all the fibers (see[30]). This is classically done by endowing F_S with a topological structure that glues together the fibers (which are computed as algebraic limits in the category of sets). One of the main difficulties here resides in the fact that the inductive limit in Cu is not the algebraic inductive limit, even in the case of the fiber of a surjective presheaf. We illustrate this situation below with an easy example. For a semigroup S in Cu and a compact Hausdorff space X , we shall

denote by $\text{Lsc}(X, S)$ the set of those functions $f: X \rightarrow S$ such that $\{t \in X \mid f(t) \gg s\}$ is open in X for all $s \in S$. If X is finite dimensional, it was shown in [3] that $\text{Lsc}(X, S) \in \text{Cu}$.

Example 2.1. Let $A = C([0, 1], M_n(\mathbb{C}))$, where $n \geq 2$. We know that $\text{Cu}(A) \cong \text{Lsc}([0, 1], \overline{\mathbb{N}})$, where $\overline{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$ (see, e.g. [21]).

Now, let $\{U_m = [\frac{1}{2} - \frac{1}{m}, \frac{1}{2} + \frac{1}{m}]\}_{m \geq 2}$, which is a sequence of decreasing closed subsets of $[0, 1]$ whose intersection is $\{1/2\}$. It is easy to verify that, in Cu ,

$$\lim_{\rightarrow} \text{Cu}(A(U_n)) = \lim_{\rightarrow} \text{Lsc}(U_n, \overline{\mathbb{N}}) = \text{Cu}(A(1/2)) = \overline{\mathbb{N}}.$$

However, the computation of the direct limit above in the category of semigroups yields $\{(a, b, c) \in \mathbb{N}^3 \mid b \leq a, c\}$.

For a surjective continuous presheaf $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$, let $F_{\mathcal{S}} := \sqcup_{x \in X} S_x$, where $S_x = \lim_{x \in \mathring{V}} S(V)$, and define $\pi: F_{\mathcal{S}} \rightarrow X$ by $\pi(s) = x$ if $s \in S_x$.

We define a *section* of $F_{\mathcal{S}}$ as a function $f: X \rightarrow F_{\mathcal{S}}$ such that $f(x) \in S_x$. We equip the set of sections with pointwise addition and order, so this set becomes an ordered semigroup. Notice also that the set of sections is closed under pointwise suprema of increasing sequences.

Any element $s \in S$ induces a section \hat{s} , which is defined by $\hat{s}(x) = \pi_x(s) \in S_x$ and will be referred to as the *section induced* by s .

Lemma 2.2. *Let $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ be a presheaf on X , and let $s, r \in S$.*

- (i) *If $\hat{s}(x) \leq \hat{r}(x)$ for some $x \in X$ then, for each $s' \ll s$ in S there is a closed set V with $x \in \mathring{V}$ such that $\pi_V(s') \leq \pi_V(r)$. In particular, $\hat{s}'(y) \leq \hat{r}(y)$ for all $y \in V$.*
- (ii) *If, further, \mathcal{S} is a sheaf, U is a closed subset of X , and $\hat{s}(x) \leq \hat{r}(x)$ for all $x \in U$, then for each $s' \ll s$ there is a closed set W of X with $U \subset \mathring{W}$ and $\pi_W(s') \leq \pi_W(r)$.*

Proof. (i): Recall that $S_x = \lim S(V_n)$, where (V_n) is a decreasing sequence of closed sets whose intersection is x (we may take $V_1 = X$, and $x \in \mathring{V}_n$ for all n). Then, by Proposition 1.3, two elements s, r satisfy $\hat{s}(x) = \pi_x(s) \leq \pi_x(r) = \hat{r}(x)$ in S_x if and only if for all $s' \ll s$ there exists $j \geq 1$ such that $\pi_{V_j}(s') \leq \pi_{V_j}(r)$, and in particular $\hat{s}'(y) \leq \hat{r}(y)$ for all y in V_j .

(ii): Assume now that \mathcal{S} is a sheaf, and take $s' \ll s$. Apply (i) to each $x \in U$, so that we can find U_x with $x \in \mathring{U}_x$ such that $\pi_{U_x}(s') \leq \pi_{U_x}(r)$. By compactness of U , there are a finite number U_{x_1}, \dots, U_{x_n} whose interiors cover U . Put $W = \cup_i U_{x_i}$. As \mathcal{S} is a sheaf and $\pi_{U_{x_i}}(s') \leq \pi_{U_{x_i}}(r)$ for all i , it follows that $\pi_W(s') \leq \pi_W(r)$. \square

Following Lemma 2.2, our aim is to define a topology in $F_{\mathcal{S}}$ for which the induced sections will be continuous. Instead of abstractly considering the final topology generated by the induced sections, we define a particular topology which will satisfy our needs. Given U an open set in X and $s \in S$, put

$$U_s^{\gg} = \{a_x \in F_{\mathcal{S}} \mid \hat{s}'(x) \ll a_x \text{ for some } x \in U \text{ and some } s \ll s'\},$$

and equip $F_{\mathcal{S}}$ with the topology generated by these sets.

Now consider an induced section \hat{s} for some $s \in S$, and an open set of the form U_r^{\gg} for some $r \in S$ and $U \subseteq X$. Suppose $x \in \hat{s}^{-1}(U_r^{\gg})$. Note that $x \in U$ and that $\hat{s}(x) \gg \hat{s}'(x)$ for some $s' \gg r$. Using that $s' = \sup(s'_n)$ for a rapidly increasing sequence (s'_n) , there exists n_0

such that $r \ll s'_{n_0} \ll s'$. Hence, by Lemma 2.2, there is a closed set V such that $x \in \mathring{V}$ and $\hat{s}'_{n_0}(y) \ll \hat{s}(y)$ for all y in V . Thus, $x \in U \cap \mathring{V} \subseteq \hat{s}^{-1}(U_r^{\gg})$, proving that $\hat{s}^{-1}(U_r^{\gg})$ is open in X , from which it easily follows that \hat{s} is continuous with this topology.

Remark 2.3. Notice that if S is a surjective presheaf, then any element $a \in S_x$ can be written as $a = \sup(\hat{s}_n(x))$, where s_n is a rapidly increasing sequence in S . This is possible as the map $S \rightarrow S_x$ is surjective, hence $a = \pi_x(s)$ for some $s \in S$, and $s = \sup s_n$ for such a sequence.

The following result gives another characterization of continuity that will prove useful in the sequel.

Proposition 2.4. *Let X be a compact Hausdorff space and S be a continuous surjective presheaf on Cu . Then, for a section $f: X \rightarrow F_S$, the following conditions are equivalent:*

- (i) f is continuous.
- (ii) For all $x \in X$ and $a_x \in S_x$ such that $a_x \ll f(x)$, there exist a closed set V with $x \in \mathring{V}$ and $s \in S$ such that $\hat{s}(x) \gg a_x$ and $\hat{s}(y) \ll f(y)$ for all $y \in V$.

Proof. Let $f: X \rightarrow F_S$ be a section satisfying (ii) and consider an open set of the form U_r^{\gg} for some open set $U \subseteq X$ and $r \in S$. Then

$$\begin{aligned} f^{-1}(U_r^{\gg}) &= \{y \in X \mid f(y) \gg \hat{r}'(y) \text{ for some } y \in U \text{ and for some } r' \gg r\} \\ &= \{y \in U \mid f(y) \gg \hat{r}'(y) \text{ for some } r' \gg r\}. \end{aligned}$$

For each y in the above set there exists $r' \gg r$ such that $\hat{r}'(y) \ll f(y)$. Using property (ii) there exists $s \in S$ such that $\hat{r}'(y) \ll \hat{s}(y) \ll f(y)$ and $\hat{s}(x) \ll f(x)$ for all $x \in \mathring{V}$ where V is a closed set of X . Furthermore, we can find $r'' \in S$ such that $r \ll r'' \ll r'$, and use Lemma 2.2 to conclude that $\hat{r}''(z) \ll \hat{s}(z) \ll f(z)$ for all z in an open set $W \subseteq X$, proving that $f^{-1}(U_r^{\gg})$ is open. Therefore f is continuous.

Now, let $f: X \rightarrow F_S$ be continuous, $x \in X$ and $a_x \ll f(x)$. Using Remark 2.3, we can write $f(x) = \sup(\hat{s}_n(x))$ where (s_n) is a rapidly increasing sequence in S , and hence we can find $s \ll s' \in S$ such that

$$a_x \ll \hat{s}(x) \ll \hat{s}'(x) \ll f(x),$$

where $s, s' \in S$.

Let U be any open neighborhood of x , and consider the open set $f^{-1}(U_s^{\gg})$. Note that it contains x and that for any $z \in f^{-1}(U_s^{\gg})$, we have $f(z) \gg \hat{t}(z)$ for some $t \gg s$. Hence, for any closed set V contained in $f^{-1}(U_s^{\gg})$ such that $x \in \mathring{V}$, we have $f(z) \gg \hat{s}(z)$ for all $z \in V$. Thus, condition (ii) holds. \square

Let X be a compact Hausdorff space and let S be a continuous presheaf on Cu . We will denote the set of continuous sections of the space F_S by $\Gamma(X, F_S)$, which is equipped with pointwise order and addition. Notice that there is an order-embedding

$$\Gamma(X, F_S) \rightarrow \prod_{x \in X} S_x$$

(given by $f \mapsto (f(x))_{x \in X}$).

Definition 2.5. Let X be a compact Hausdorff space. We say that a $C(X)$ -algebra A has no K_1 obstructions provided that, for all $x \in X$, the fiber A_x has stable rank one and $K_1(I) = 0$ for any closed two-sided ideal of A_x .

The class just defined was already considered, although not quite with this terminology, in [3], where various aspects of the Cuntz semigroup of these algebras were examined. We combine some of the ideas from [3] to prove the results below, which are a first step towards the computation of the Cuntz semigroup of $C(X)$ -algebras without K_1 obstructions.

Theorem 2.6. Let X be a one dimensional compact Hausdorff space and A be a $C(X)$ -algebra without K_1 obstructions. Then, the map

$$\alpha: \text{Cu}(A) \rightarrow \prod_{x \in X} \text{Cu}(A_x),$$

given by $\alpha[a] = ([a(x)])_{x \in X}$ is an order embedding. In particular, α defines an order embedding

$$\text{Cu}(A) \rightarrow \Gamma(X, F_{\text{Cu}(A)}).$$

Proof. By our assumptions on A and its fibers, we may assume that A is stable.

Let $0 < \epsilon < 1$ be fixed, and let us suppose that $a, b \in A$ are positive contractions such that $a(x) \preceq b(x)$ for all $x \in X$. Then, by the definition of the Cuntz order, since A_x is a quotient of A for each $x \in X$, there exists $d_x \in A$ such that

$$\|a(x) - d_x(x)b(x)d_x^*(x)\| < \epsilon.$$

By upper semicontinuity of the norm, the above inequality also holds in a neighborhood of x . Hence, since X is a compact set, there exists a finite cover of X , say $\{U_i\}_{i=1}^n$, and elements $(d_i)_{i=1}^n \in A$ such that $\|a(x) - d_i(x)b(x)d_i^*(x)\| < \epsilon$, for all $x \in \overline{U_i}$ and $1 \leq i \leq n$. As X is one dimensional, we may assume that $\{U_i\}$ and $\{\overline{U_i}\}$ have order at most two.

Choose, by Urysohn's Lemma, functions λ_i that are 1 in the closed sets $U_i \setminus (\bigcup_{j \neq i} U_j)$ and 0 in U_i^c . Using these functions we define $d(x) = \sum_{i=1}^n \lambda_i(x)d_i(x)$. Set $V = X \setminus (\bigcup_{i \neq j} (U_i \cap U_j))$ which is a closed set, and it is easy to check that d satisfies

$$(1) \quad \|a(x) - d(x)b(x)d^*(x)\| < \epsilon$$

for all $x \in V$.

Again, choose for $i < j$ functions $\alpha_{i,j}$ such that $\alpha_{i,j}$ is one on $\overline{U_i} \cap \overline{U_j}$ and zero on $\overline{U_k} \cap \overline{U_l}$ whenever $\{k, l\} \neq \{i, j\}$. We define $c(x) = \sum_{i < j} \alpha_{i,j}(x)d_i(x)$, put $U = (\bigcup_{i \neq j} (U_i \cap U_j))^c = V^c$ and notice that c satisfies

$$(2) \quad \|a(x) - c(x)b(x)c^*(x)\| < \epsilon$$

for all $x \in \overline{U}$.

Now, by [16, Lemma 2.2], equations (1) and (2), and taking into account that the norm of an element is computed fiberwise ([6]), we have that

$$\pi_V((a - \epsilon)_+) \preceq \pi_V(b) \text{ and } \pi_{\overline{U}}((a - \epsilon)_+) \preceq \pi_{\overline{U}}(b).$$

Therefore

$$([\pi_V(a - \epsilon)_+], [\pi_{\overline{U}}(a - \epsilon)_+]) \leq ([\pi_V(b)], [\pi_{\overline{U}}(b)])$$

in the pullback semigroup $\text{Cu}(A(V)) \oplus_{\text{Cu}(A(\overline{U \cap V}))} \text{Cu}(A(\overline{U}))$. Since A can also be written as the pullback $A = A(V) \oplus_{A(\overline{U \cap V})} A(\overline{U})$ along the natural restriction maps (see [12, Lemma 2.4], and also [15, Proposition 10.1.13]), we can apply [3, Theorem 3.2], to conclude that $(a - \epsilon)_+ \preceq b$. Thus $a \preceq b$, and the result follows. \square

Corollary 2.7. *Let X be a one dimensional compact Hausdorff space, and let A be a $C(X)$ -algebra without K_1 obstructions. Then, $\text{Cu}_A: \mathcal{V}_X \rightarrow \text{Cu}, U \mapsto \text{Cu}(A(U))$, is a surjective continuous sheaf.*

Proof. We know already that Cu_A is a surjective continuous presheaf. Let U and $V \in \mathcal{V}_X$ be such that $U \cap V \in \mathcal{V}_X$. Let $W = U \cup V$. We know then that $A(W)$ is isomorphic to the pullback $A(U) \oplus_{A(U \cap V)} A(V)$. Since $A(W)$ is a $C(W)$ -algebra without K_1 obstructions, we may apply Theorem 2.6 to conclude that the map $\text{Cu}(A(W)) \rightarrow \prod_{x \in W} \text{Cu}(A_x)$ (given by $[a] \mapsto ([a(x)])$) is an order-embedding. Then [3, Theorem 3.3] implies that the natural map $\text{Cu}(A(W)) \rightarrow \text{Cu}(A(U)) \oplus_{\text{Cu}(A(U \cap V))} \text{Cu}(A(V))$ is surjective. Since it is also an order-embedding, by [3, Theorem 3.2], we obtain that it is an isomorphism. \square

3. PIECEWISE CHARACTERISTIC SECTIONS

In this section we will show that, under additional assumptions, the map in Theorem 2.6 is also surjective, proving that there exists an isomorphism in the category Cu between $\text{Cu}(A)$ and $\Gamma(X, F_{\text{Cu}(A)})$.

Recall that, if $s \ll r \in S$, then $\pi_x(s) = \hat{s}(x) \ll \hat{r}(x) = \pi_x(r)$ for all x . This comes from the fact that the induced maps belong to the category Cu , and so they preserve the compact containment relation. We continue to assume throughout that X is a compact Hausdorff space, which is also second countable. We shall use below $\partial(U)$ to denote the *boundary* of a set U , that is, $\partial(U) = \overline{U} \setminus \overset{\circ}{U}$.

Lemma 3.1. *Let $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ be a surjective presheaf of semigroups on X .*

(i) *Let $f, g \in \Gamma(X, F_{\mathcal{S}})$, and V a closed subset of X such that $f(y) \ll g(y)$ for all $y \in V$. Put*

$$g_{V,f}(x) = \begin{cases} g(x) & \text{if } x \notin V \\ f(x) & \text{if } x \in V \end{cases}$$

Then $g_{V,f} \in \Gamma(X, F_{\mathcal{S}})$.

(ii) *If $g \in \Gamma(X, F_{\mathcal{S}})$ and $x \in X$, there exist a decreasing sequence (V_n) of closed sets (with $x \in \overset{\circ}{V}_n$ for all n) and a rapidly increasing sequence (s_n) in S such that $g = \sup_n g_{V_n, s_n}$.*

Proof. (i): Using the fact that both f and g are continuous, it is enough to check that condition (ii) in Proposition 2.4 is verified for $x \in \partial(V)$. Thus, let a_x be such that $a_x \ll g_{V,f}(x) = f(x) \ll g(x)$. By continuity of f , there is a closed subset U with $x \in \overset{\circ}{U}$ and $s \in S$ such that $a_x \ll \hat{s}(x)$ and $\hat{s}(y) \ll f(y)$ for all $y \in U$. As s is a supremum of a rapidly increasing sequence, we may find $s' \ll s$ with $a_x \ll \hat{s}'(x)$.

Next, as g is also continuous, there are $t \in S$ and a closed set U' with $x \in \overset{\circ}{U}'$ such that $f(x) \ll \hat{t}(x)$ and $\hat{t}(y) \ll g(y)$ for all $y \in U'$. Since $\hat{s}(x) \ll \hat{t}(x)$ and $s' \ll s$, we now use Lemma 2.2 to find W with $\hat{s}'(y) \ll \hat{t}(y)$ for all $y \in W$. Now condition (ii) in Proposition 2.4 is verified using the induced section s' and the closed set $U \cap U' \cap W$.

(ii): Write $g(x) = \sup_n \hat{s}_n(x)$, where (s_n) is a rapidly increasing sequence in S (see Remark 2.3).

Since $s_1 \ll s_2$ and g is continuous, condition (ii) of Proposition 2.4 applied to $\hat{s}_2(x) \ll g(x)$ yields $t \in S$ and a closed set U_1 whose interior contains x such that $\hat{s}_2(x) \ll \hat{t}(x)$ and $\hat{t}(y) \ll g(y)$ for all $y \in V_1$. We now apply Lemma 2.2, so that there is another closed set U'_1 (with $x \in \overset{\circ}{U}'_1$) so that $\hat{s}_1(y) \ll \hat{t}(y)$ for any $y \in U'_1$. Let $V_1 = U_1 \cap U'_1$ and for each $y \in V_1$, we have $\hat{s}_1(y) \ll \hat{t}(y) \ll g(y)$. Continue in this way with the rest of the s_n 's, and notice that we can choose the sequence (V_n) in such a way that $\bigcap V_n = \{x\}$. \square

Using the previous lemma we can describe compact containment in $\Gamma(X, F_{\text{Cu}(A)})$.

Proposition 3.2. *Let $S: \mathcal{V}_X \rightarrow \text{Cu}$ be a surjective presheaf of semigroups on X . For f, g in $\Gamma(X, F_{\text{Cu}(A)})$, the following statements are equivalent:*

(i) $f \ll g$.

(ii) *For all $x \in X$ there exists a_x with $f(x) \ll a_x \ll g(x)$ and such that if $s \in S$ satisfies $a_x \ll \hat{s}(x)$ and $\hat{s}(y) \ll g(y)$ for y in a closed set U whose interior contains x , then there exists a closed set $V \subseteq U$ with $x \in \overset{\circ}{V}$ and $f(y) \leq \hat{s}(y) \leq g(y)$ for all $y \in V$.*

Proof. (i) \implies (ii): Given $x \in X$, use Lemma 3.1 to write $g = \sup_n g_{V_n, s_n}$, where (s_n) is rapidly increasing in S and (V_n) is a decreasing sequence of closed sets whose interior contain x . Since $f \ll g$, there is n such that

$$f \leq g_{V_n, s_n} \leq g_{V_{n+1}, s_{n+1}} \leq g.$$

Let $a_x = g_{V_{n+1}, s_{n+1}}(x) = \hat{s}_{n+1}(x)$, which clearly satisfies $f(x) \leq \hat{s}_n(x) \ll \hat{s}_{n+1}(x) \ll g(x)$. Assume now that $s \in S$ and U is a closed set with $x \in \overset{\circ}{U}$ such that $a_x \ll \hat{s}(x)$ and $\hat{s}(y) \ll g(y)$ for all $y \in U$. Since $s_n \ll s_{n+1}$ and $\hat{s}_{n+1}(x) \ll \hat{s}(x)$, there is by Lemma 2.2 a closed set V with $x \in \overset{\circ}{V}$ (and we may assume $V \subset V_{n+1} \cap U$) such that $\hat{s}_n(y) \leq \hat{s}(y)$ for all $y \in V$. Thus $f(y) \leq \hat{s}_n(y) \leq \hat{s}(y) \leq g(y)$ for all $y \in V$.

(ii) \implies (i): Suppose now that $g \leq \sup(g_n)$, where (g_n) is an increasing sequence in $\Gamma(X, F_S)$. Let $x \in X$, and write $g = \sup g_{V_n, s_n}$ as in Lemma 3.1, where (s_n) is a rapidly increasing sequence in S . Our assumption provides us first with a_x such that $f(x) \ll a_x \ll g(x)$. In particular, there is m such that $a_x \ll \hat{s}_m(x) \ll \hat{s}_{m+1}(x) \ll \hat{s}_{m+2}(x) \ll g(x)$, and hence there exists k with $\hat{s}_{m+1}(x) \ll g_k(x)$.

As g_k is continuous, condition (ii) in Proposition 2.4 implies that we may find $s \in S$ and a closed set U with $x \in \overset{\circ}{U}$ such that $\hat{s}_{m+1}(x) \ll \hat{s}(x)$ and $\hat{s}(y) \ll g_k(y)$ for all $y \in U$. Now, as $s_m \ll s_{m+1}$, there exists a closed subset V with $x \in \overset{\circ}{V}$ and $\hat{s}_m(y) \leq \hat{s}(y)$ for all $y \in V$, whence $\hat{s}_m(y) \leq g(y)$ for all $y \in U \cap V$.

Since also $\hat{s}_m(y) \ll g(y)$ for all $y \in V_m$, there is by assumption a closed set $W \subseteq V_m \cap V$ (whose interior contains x) such that $f(y) \leq \hat{s}_m(y) \leq g_k(y)$ for all $y \in W$. Now, by a standard compactness argument we may choose l such that $f \leq g_l$. \square

Lemma 3.3. *Let $S: \mathcal{V}_X \rightarrow \text{Cu}$ be a surjective presheaf of semigroups. Then, the morphism*

$$\begin{aligned} \alpha: S &\rightarrow \Gamma(X, F_S) \\ s &\mapsto \hat{s} \end{aligned}$$

preserves compact containment and suprema.

Proof. Using condition (ii) of Proposition 2.4 it is easy to verify that, if (f_n) is an increasing sequence in $\Gamma(X, F_S)$, then its pointwise supremum is also a continuous section.

Assume now that $s \ll r$ in S . Write $r = \sup(r_n)$, where (r_n) is a rapidly increasing sequence in S . We may find m such that

$$s \ll r_m \ll r.$$

Take $a_x = \hat{r}_m(x)$. Suppose that $t \in S$ satisfies $a_x \ll \hat{t}(x)$ and $\hat{t}(y) \ll \hat{r}(y)$ for y in a closed subset U whose interior contains x . By Lemma 2.2, applied to $\hat{r}_m(x) \leq \hat{t}(x)$ and $s \ll r_m$, there is a closed set V such that $x \in \overset{\circ}{V}$ and $\hat{s}(y) \leq \hat{t}(y)$ for $y \in V$. Thus, for any $y \in V \cap U$, we have $\hat{s}(y) \leq \hat{t}(y) \leq \hat{r}(y)$. This verifies condition (ii) in Proposition 3.2, whence $\hat{s} \ll \hat{r}$. \square

Corollary 3.4. *Let $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ be a surjective sheaf of semigroups on X , $f \in \Gamma(X, F_S)$, $s \in S$, and let V be a closed subset of X . If $\hat{s}(x) \leq f(x)$ for all $x \in V$ and $s' \ll s$, then there is a closed subset W of X with $V \subset \overset{\circ}{W}$ such that $\pi_W(s') \ll f|_W$.*

Proof. Let $s' \ll t' \ll t \ll s$ in S . For each $x \in V$, there is by Proposition 2.4 a closed set U_x whose interior contains x , and $r_x \in S$ such that $\hat{t}(x) \ll \hat{r}_x(x)$, and $\hat{r}_x(y) \leq f(y)$ for all $y \in U_x$. Now apply condition (i) of Lemma 2.2 to $t' \ll t$ in order to find another closed set V_x such that $x \in \overset{\circ}{V}_x$ and $\hat{t}'(y) \leq \hat{r}_x(y)$ for $y \in V_x$. Letting $W_x = U_x \cap V_x$, we have $\hat{t}'(y) \ll f(y)$ for all $y \in W_x$. Since $V \subseteq \bigcup_x \overset{\circ}{W}_x$, and V is closed, we may find a finite number of W_x 's that cover V , whose union is the closed set W we are after. Since \mathcal{S} is a sheaf, it follows that $\pi_W(t') \leq f|_W$, and by Lemma 3.3 we see that $\pi_W(s') \ll \pi_W(t') \leq f|_W$, as desired. \square

We now proceed to define a class of continuous sections that will play an important role. This will be a version, for presheaves on spaces of dimension one, of the notion of piecewise characteristic function given in [3, Definitions 2.4 and 5.9]. We show below that, for a surjective sheaf of semigroups $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ on a one dimensional space X , every element in $\Gamma(X, F_S)$ can be written as the supremum of a rapidly increasing sequence of piecewise characteristic sections. From this, we can conclude that $\Gamma(X, F_S)$ is an object in Cu . Just as in [3], we could define piecewise characteristic sections for spaces of arbitrary (finite) dimension and make the case that $\Gamma(X, F_S)$ belongs to Cu as well. This is however technically much more involved and beyond the scope of this paper, whence it will be pursued elsewhere.

Definition 3.5. (*Piecewise characteristic sections*) *Let X be a one dimensional compact Hausdorff space. Let $\{U_i\}_{i=1 \dots n}$ be an open cover of X such that the order of $\{U_i\}$ and $\{\bar{U}_i\}$ is at most two. Assume also that $\dim(\partial(\bar{U}_i)) = 0$ for all i .*

Let $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ be a presheaf of semigroups on X . For $i \in \{1, \dots, n\}$, choose elements $s_i \in S$ and $s_{\{i,j\}} \in S$ whenever $i \neq j$, such that

$$\hat{s}_i(x) \leq \hat{s}_{\{i,j\}}(x) \text{ for all } x \text{ in } \overline{\partial(U_i \cap U_j)} \cap U_i.$$

We define a piecewise characteristic section as

$$g(x) = \begin{cases} \hat{s}_i(x) & \text{if } x \in U_i \setminus (\bigcup_{j \neq i} U_j) \\ \hat{s}_{\{i,j\}}(x) & \text{if } x \in U_i \cap U_j \end{cases}.$$

By an argument similar to the one in Lemma 3.1, it follows that piecewise characteristic sections are continuous.

Remark 3.6. In the case of zero dimensional spaces, piecewise characteristic sections are much easier to define. Given an open cover $\{U_i\}_{i=1,\dots,n}$ consisting of pairwise disjoint clopen sets, a presheaf of semigroups \mathcal{S} on Cu and elements $s_1, \dots, s_n \in \mathcal{S}$, a piecewise characteristic section in this setting is an element $g \in \Gamma(X, F_{\mathcal{S}})$ such that $g(x) = \hat{s}_i(x)$, whenever $x \in U_i$.

If $f \in \Gamma(X, F_{\mathcal{S}})$ and g is a piecewise characteristic section such that $g \ll f$, then we say that g is a piecewise characteristic section of f and we will denote the set of these sections by $\chi(f)$.

Lemma 3.7. *Let X be a one dimensional compact Hausdorff space, and $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ a surjective presheaf of semigroups. If $f \in \Gamma(X, F_{\mathcal{S}})$, then*

$$f = \sup\{g \mid g \in \chi(f)\}.$$

Proof. Let $x \in X$. By Lemma 3.1, we may write $f = \sup f_{V_n, s_n}$, where (V_n) is a decreasing sequence of closed sets with $x \in \overset{\circ}{V}_n$ and (s_n) is rapidly increasing. By construction, $f_{V_n, s_n}(y) = \hat{s}_n(y) \ll f(y)$ for all $y \in V_n$.

Now define

$$h_n(y) = \begin{cases} \hat{s}_n(y) & \text{if } y \in \overset{\circ}{V}_n \\ 0 & \text{otherwise} \end{cases}.$$

It is easy to verify, using Proposition 3.2, that $h_n \ll f$, and also that each h_n is a piecewise characteristic section for f . Using this fact for all $x \in X$, we conclude that $f = \sup\{g \mid g \in \chi(f)\}$. \square

Proposition 3.8. *Let X be a compact Hausdorff space with $\dim(X) \leq 1$, and let $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ be a surjective sheaf of semigroups. Suppose $h_1, h_2, f \in \Gamma(X, F_{\mathcal{S}})$ such that $h_1, h_2 \ll f$. Then, there exists $g \in \chi(f)$ such that $h_1, h_2 \ll g$. In particular, $\chi(f)$ is an upwards directed set.*

Proof. Assume first that X has dimension 0. Writing f as in condition (ii) of Lemma 3.1 we can find, for each $x \in X$, an open set V_x that contains x , and elements $s'_x \ll s_x \ll s''_x \in \mathcal{S}$ such that

$$(3) \quad h_1(y), h_2(y) \ll \hat{s}'_x(y) \ll \hat{s}_x(y) \ll \hat{s}''_x(y) \ll f(y) \quad \text{for all } y \in \overline{V}_x.$$

Using compactness and the fact that X is zero dimensional, there are $x_1, \dots, x_n \in X$ and (pairwise disjoint) clopen sets $\{V_i\}_{i=1,\dots,n}$ with $V_i \subseteq V_{x_i}$ and such that $X = \cup_i V_i$. Put $s_i = s_{x_i}$, $s'_i = s'_{x_i}$ and $s''_i = s''_{x_i}$. Define, using this cover, a piecewise characteristic section g as $g(x) = \hat{s}_i(x)$ if $x \in V_i$. It now follows from (3) that $h_1, h_2 \ll g \ll f$ (the elements s'_i, s''_i are used here to obtain compact containment).

We turn now to the case where X has dimension 1, and start as in the previous paragraph, with some additional care. Choose, for each x , a δ_x -ball V_x'' (where $\delta_x > 0$) centered at x and elements $s'_x \ll s_x \ll s''_x$ such that condition (3) is satisfied (for all $y \in \overline{V}_x''$). Denote by $V_x' \subseteq V_x''$ the cover consisting of $\delta_x/2$ -balls. By compactness we obtain a finite cover $\{V'_{x_1}, \dots, V'_{x_n}\}$. Using [20, Lemma 8.1.1] together with the fact that X has dimension 1, this cover has a refinement $\{V_i\}_{i=1}^n$ such that $\{V_i\}$ and $\{\overline{V}_i\}$ have both order at most 2 and such that $\partial(V_i)$ has dimension 0 for each i . As before, set $s_i = s_{x_i}$, $s'_i = s'_{x_i}$ and $s''_i = s''_{x_i}$.

Let Y be the closed set $\cup_i \partial(V_i)$, which also has dimension 0. Put $\delta = \min\{\delta_{x_i}/3\}$. By construction, there is a δ -neighborhood V_i^δ such that $V_i^\delta \subseteq V_i''$. As in the proof of Lemma 3.7, we see that the sections

$$g_i(y) = \begin{cases} \hat{s}_i''(y) & \text{if } y \in V_i^\delta \\ 0 & \text{otherwise} \end{cases}$$

satisfy $g_i \ll f$. We now restrict to Y and proceed as in the argument of the zero dimensional case above. In this way, we obtain piecewise characteristic sections $g_Y, g_Y', g_Y'' \in \Gamma(Y, F_S)$, defined by some open cover $\{W_i\}_{i=1}^m$ (of pairwise disjoint clopen sets of Y) and elements $t_i \ll t_i' \ll t_i'' \in S$ in such a way that $g_Y(y) = \hat{t}_i(y)$, $g_Y'(y) = \hat{t}_i'(y)$ and $g_Y''(y) = \hat{t}_i''(y)$ whenever $y \in W_i$, and such that

$$(4) \quad \pi_Y(g_i) \ll g_Y \ll g_Y' \ll g_Y'' \ll \pi_Y(f) \text{ for all } i = 1, \dots, n.$$

Observe that we can choose the W_i of arbitrarily small size, thus in particular we may assume that each one is contained in a $\delta/6$ -ball. In this way, whenever $\overline{W}_i \cap \overline{W}_j \neq \emptyset$, we have $W_i \subseteq V_j^\delta$. Therefore, if $x \in W_i$, it follows from (4) that

$$\hat{s}_j''(x) = g_j(x) \leq g_Y(x) = \hat{t}_i(x).$$

By condition (ii) in Lemma 2.2, applied to the previous inequality, there is $\epsilon > 0$ such that $\hat{s}_j(x) \leq \hat{t}_i(x)$ for all $x \in W_i^\epsilon$. Since the W_i are pairwise disjoint clopen sets, we can choose ϵ such that the sets W_i^ϵ are still pairwise disjoint. Further, since also $\hat{t}_i''(y) \leq f(y)$ for $y \in W_i$ and $t_i' \ll t_i''$, we may apply Corollary 3.4 to obtain $\pi_{\overline{W}_i^\epsilon}(t_i') \ll \pi_{\overline{W}_i^\epsilon}(f)$ (further decreasing ϵ if necessary). As for each i , we can find U_i with $W_i^{\epsilon/2} \subseteq U_i \subseteq W_i^\epsilon$ with zero dimensional boundary, after a slight abuse of notation we shall assume that W_i^ϵ itself has zero dimensional boundary. Put $Y^\epsilon = \cup_{i=1}^m W_i^\epsilon$. Notice now that, for $i, k < l$, the closed sets $V_i \setminus (Y^\epsilon \cup \cup_{j \neq i} V_j)$ and $(V_k \cap V_l) \setminus Y^\epsilon$ are also pairwise disjoint, whence they admit pairwise disjoint ϵ' -neighborhoods (for a sufficiently small ϵ'). As before, we shall also assume these neighborhoods have zero dimensional boundary.

Now consider the cover that consists of the sets

$$\{W_i^\epsilon, i = 1, \dots, m, (V_i \setminus (Y^\epsilon \cup \cup_{j \neq i} V_j))^{\epsilon'}, i = 1, \dots, n, (V_k \cap V_l \setminus (Y^\epsilon))^{\epsilon'}, k < l\},$$

and define a piecewise characteristic section g as follows

$$g(x) = \begin{cases} \hat{t}_i(x) & \text{if } x \in W_i^\epsilon \\ \hat{s}_i(x) & \text{if } x \in (V_i \setminus (Y^\epsilon \cup \cup_{j \neq i} V_j))^{\epsilon'} \setminus Y^\epsilon \\ \hat{s}_k(x) & \text{if } x \in (V_k \cap V_l \setminus (Y^\epsilon))^{\epsilon'} \setminus Y^\epsilon \text{ for } k < l. \end{cases}$$

That $h_1, h_2 \leq g$ follows by construction of g . It remains to show that $g \ll f$. This also follows from our construction, using condition (ii) of Proposition 3.2. For example, given $x \in W_i^\epsilon$, we have

$$g(x) = \hat{t}_i(x) \ll \hat{t}_i'(x) \ll f(x).$$

If now $s \in S$ satisfies that $\hat{t}_i'(x) \ll \hat{s}(x)$ and $\hat{s}(y) \ll f(y)$ for y in a closed set (whose interior contains x) then, since $t_i \ll t_i'$, we may find (again by Lemma 2.2) a smaller closed set (contained in W_i^ϵ and with interior containing x) such that $g(y) = \hat{t}_i(y) \leq \hat{s}(y) \leq f(y)$ for y in that set. \square

Proposition 3.9. *Let X be a one dimensional compact Hausdorff space, and let $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ be a surjective sheaf of semigroups with S countably based. If $f \in \Gamma(X, F_S)$, then f is the supremum of a rapidly increasing sequence of elements from $\chi(f)$.*

Proof. Let us define a new topology on F_S . Let $s \in S$ and let U be an open set in X . Consider the topology generated by the sets

$$U_s^{\ll} = \{y \in F_S \mid \hat{s}(x) \gg y \text{ for some } x \in U\}.$$

We claim that, under this topology, F_S is second countable. Let $\{U_n\}$ be a basis of X , and $\{s_n\}_{n \in \mathbb{N}}$ be a dense subset of S . Therefore the collections of sets $\{(U_n)_{s_i}^{\ll}\}_{n,i \in \mathbb{N}}$ is a countable basis for F_S . Indeed, given an open set U of X and $s \in S$, find sequences (U_{n_i}) and (s_{m_j}) such that $U = \cup U_{n_i}$ and $s = \sup s_{m_j}$. Then

$$U_s^{\ll} = \cup (U_{n_i})_{s_{m_j}}^{\ll}.$$

Now, for $f \in \Gamma(X, F_S)$, put $U_f = \{a_x \in F_S \mid a_x \ll f(x) \text{ for } x \in U\}$. This set is open in the topology we just have defined. To see this, let $a_x \in U_f$, and invoke Proposition 2.4 to find an open set V and $s \in S$ such that $a_x \ll \hat{s}(x)$ and $\hat{s}(y) \ll f(y)$ for all $y \in V$. It then follows that $a_x \in V_s^{\ll} \subseteq U_f$.

Using Lemma 3.7, we see that $U_f = \cup_{g \in \chi(f)} U_g$. Since F_S is second countable, it has the Lindelöf property, whence we may find a sequence (g_n) in $\chi(f)$ such that $U_f = \cup_n U_{g_n}$. This sequence may be taken to be increasing by Proposition 3.8. Translating this back to $\Gamma(X, F_S)$, we get $f = \sup(g_n)$. \square

Assembling our observations we obtain the following:

Theorem 3.10. *Let X be a one dimensional compact Hausdorff space, and let $\mathcal{S}: \mathcal{V}_X \rightarrow \text{Cu}$ be a surjective sheaf of semigroups such that S is countably based. Then, the semigroup $\Gamma(X, F_S)$ of continuous sections belongs to the category Cu .*

The next result shows, in a particular case, the existence of an induced section between any two compactly contained piecewise characteristic sections.

Proposition 3.11. *Let X be a one dimensional compact Hausdorff space and let A be a stable continuous field over X with no K_1 obstructions. Let $f \ll g$ be elements in $\Gamma(X, F_{\text{Cu}(A)})$ such that g is a piecewise characteristic section. Then there exists an element $h \in A$ which satisfies $f(x) \leq [\pi_x(h)] \leq g(x)$ for all $x \in X$.*

Proof. Since g is a piecewise characteristic section there is a cover $\{U_i\}_{i=1}^n$ of X such that both $\{U_i\}$ and $\{\bar{U}_i\}$ have order at most 2, and there are elements $[a_i], [a_{\{i,j\}}]$ in $\text{Cu}(A)$ which are the values that g takes (according to Definition 3.5).

For $\epsilon > 0$, let g_ϵ be the section defined on the same cover as g and that takes values $[(a_i - \epsilon)_+], [a_{\{i,j\}}]$. As $g = \sup_\epsilon g_\epsilon$ and $f \ll g$, we may choose $\epsilon > 0$ such that $f \leq g_\epsilon$, and in particular

$$f(x) \leq \pi_x([(a_i - \epsilon)_+]) \ll \pi_x([a_i]) \text{ for all } x \text{ in } U_i \setminus (\cup_{j \neq i} U_j).$$

Notice now that the closed sets $\overline{\partial(U_i \cap U_j) \cap U_i}$ and $\overline{\partial(U_k \cap U_l) \cap U_l}$ are pairwise disjoint whenever $(i, j) \neq (k, l)$. (This follows from elementary arguments together with the assumption that the cover $\{\bar{U}_i\}$ has order at most 2.)

Furthermore, by definition of g we have $\pi_x([a_i]) \leq \pi_x([a_{\{i,j\}}])$ for all $x \in \overline{\partial(U_i \cap U_j) \cap U_i}$. Therefore, there exists by Corollary 3.4 a neighborhood $W_{i,j}$ of $\overline{\partial(U_i \cap U_j) \cap U_i}$ for which

$$\pi_{\overline{W}_{i,j}}([a_i]) \leq \pi_{\overline{W}_{i,j}}([a_{\{i,j\}}]).$$

We may assume without loss of generality that the closures $\overline{W}_{i,j}$ are pairwise disjoint sets. Since also $\overline{\partial(U_i \cap U_j) \cap U_i} \cap \overline{U}_k = \emptyset$ whenever $k \neq i, j$, we may furthermore assume that $W_{i,j} \cap \overline{U}_k = \emptyset$ for $k \neq i, j$.

By Proposition 1.1 there exist unitaries $u_{i,j} \in \mathcal{U}(A(\overline{W}_{i,j})^\sim)$ such that

$$u_{i,j} \pi_{\overline{W}_{i,j}}((a_i - \epsilon)_+) u_{i,j}^* \in \text{Her}(\pi_{\overline{W}_{i,j}}(a_{\{i,j\}})).$$

Now, as A and $A(\overline{W}_{i,j})$ are stable, the unitary groups of their multiplier algebras are connected in the norm topology (see, e.g. [31, Corollary 16.7]). Furthermore, since the natural map $\pi_{\overline{W}_{i,j}} : A \rightarrow A(\overline{W}_{i,j})$ induces a surjective morphism $\mathcal{M}(A) \rightarrow \mathcal{M}(A(\overline{W}_{i,j}))$ (by, e.g. [31, Theorem 2.3.9]), we can find, for each unitary $u_{i,j}$, a unitary lift $\tilde{u}_{i,j}$ in $\mathcal{M}(A)$.

We now have continuous paths of unitaries $w_{i,j} : [0, 1] \rightarrow \mathcal{U}(\mathcal{M}(A))$ such that $w_{i,j}(0) = 1$ and $w_{i,j}(1) = \tilde{u}_{i,j}$. Put $\gamma = \min\{\text{dist}(\overline{W}_{i,j}, \overline{W}_{k,l} \mid (i, j) \neq (k, l))\}$. Note that $\gamma > 0$ as the sets $\overline{W}_{i,j}$ are pairwise disjoint. For $x \in X$, define a unitary in $\mathcal{M}(A)$ by

$$w_{i,j}^x = w_{i,j} \left(\frac{(\gamma - \text{dist}(x, W_{i,j}))_+}{\gamma} \right).$$

Observe that, if $x \in W_{k,l}$, then $w_{i,j}^x = \tilde{u}_{i,j}$ if $(k, l) = (i, j)$ and equals 1 otherwise. Now put

$$w_i^x = \prod_j w_{i,j}^x.$$

Since each π_x is norm decreasing and the w_i^x are defined by products and compositions of continuous functions, we obtain, using [15, Definition 10.3.1], that for each $c \in A$, the tuple $(\pi_x(w_i^x c))_{x \in X} \in \prod_{x \in X} A_x$ defines fiberwise an element in A which we denote by $w_i c$.

Now let $\{\lambda_i\}_i$ be continuous positive real-valued functions on $[0, 1]$ whose respective supports are $\{(U_i \setminus (\cup_{j \neq i} U_j)) \cup (\cup_j W_{i,j})\}_i$ and $\{\lambda_{\{i,j\}}\}_{i,j}$ with supports $\{U_i \cap U_j\}_{i,j}$. Define the following element in A

$$h = \sum_i \lambda_i w_i (a_i - \epsilon)_+ w_i^* + \sum_{i \neq j} \lambda_{\{i,j\}} a_{\{i,j\}}.$$

We now check that $[\pi_x(h)] = g_\epsilon(x)$, and this will yield the desired conclusion.

If $x \in U_i \setminus (\cup_{j \neq i} U_j)$, then $\pi_x(h) = \lambda_i(x) \pi_x(w_i (a_i - \epsilon)_+ w_i^*)$ where $\lambda_i(x) \neq 0$, and this is equivalent to $\pi_x((a_i - \epsilon)_+)$. Hence $[\pi_x(h)] = g_\epsilon(x)$.

On the other hand, if $x \in U_i \cap U_j$ for some i, j then $\lambda_{\{i,j\}}(x) \neq 0$, and

$$\pi_x(h) = \begin{cases} \lambda_i(x) \pi_x(\tilde{u}_{i,j} (a_i - \epsilon)_+ \tilde{u}_{i,j}^*) + \lambda_{\{i,j\}}(x) \pi_x(a_{\{i,j\}}) & \text{if } x \in U_i \cap U_j \cap W_{i,j} \\ \lambda_j(x) \pi_x(\tilde{u}_{j,i} (a_j - \epsilon)_+ \tilde{u}_{j,i}^*) + \lambda_{\{i,j\}}(x) \pi_x(a_{\{i,j\}}) & \text{if } x \in U_i \cap U_j \cap W_{j,i} \\ \lambda_{\{i,j\}}(x) \pi_x(a_{\{i,j\}}) & \text{if } x \in U_i \cap U_j \setminus (W_{i,j} \cup W_{j,i}). \end{cases}$$

If, for example, $x \in U_i \cap U_j \cap W_{i,j}$, then $\pi_x(\tilde{u}_{i,j} (a_i - \epsilon)_+ \tilde{u}_{i,j}^*) \in \text{Her}(\pi_x(a_{\{i,j\}}))$, and we conclude that $[\pi_x(h)] = [\pi_x(a_{\{i,j\}})] = g_\epsilon(x)$. The other cases are treated similarly. \square

This last result (together with Proposition 3.9) proves that, with some restrictions on X and A , the set of induced sections is a dense subset of $\Gamma(X, F_{\text{Cu}(A)})$, that is, every element in $\Gamma(X, F_{\text{Cu}(A)})$ is a supremum of a rapidly increasing sequence of induced sections.

Theorem 3.12. *Let X be a one dimensional compact Hausdorff space and let A be a continuous field over X without K_1 obstructions. Then, the map*

$$\begin{array}{ccc} \alpha : \text{Cu}(A) & \rightarrow & \Gamma(X, F_{\text{Cu}(A)}) \\ s & \mapsto & \hat{s} \end{array}$$

is an order isomorphism in Cu .

Proof. Let f be a continuous section in $\Gamma(X, F_{\text{Cu}(A)})$ and use Propositions 3.9 and 3.11 to write f as the supremum of a rapidly increasing sequence of induced sections $f = \sup_n \hat{s}_n$. Since α is an order embedding (by Theorem 2.6) and $\alpha(s_n) = \hat{s}_n$, the sequence s_n is also increasing in $\text{Cu}(A)$ and thus we can define $s = \sup_n s_n \in \text{Cu}(A)$. The result now follows using Lemma 3.3. \square

Since the conditions on the fibers in the previous Theorem are satisfied by simple AI-algebras we obtain the following.

Corollary 3.13. *Let X be a one dimensional compact Hausdorff space and let A be a continuous field over X such that A_x is a simple AI-algebra for all $x \in X$. Then $\text{Cu}(A) \cong \Gamma(X, F_{\text{Cu}(A)})$.*

4. THE SHEAF $\text{Cu}_A(-)$

For a compact Hausdorff space X , denote by \mathcal{C}_X the category whose objects are the $C(X)$ -algebras, and the morphisms between objects are those $*$ -homomorphisms such that commute with the (respective) structure maps.

Denote by \mathcal{S}_{Cu} the category which as objects has the presheaves $\text{Cu}_A(-)$ on X , where A belongs to \mathcal{C}_X , and the maps are presheaf homomorphisms. The following holds by definition:

Lemma 4.1. *The assignment*

$$\begin{array}{ccc} \text{Cu}_{(\cdot)} : \mathcal{C}_X & \rightarrow & \mathcal{S}_{\text{Cu}} \\ A & \mapsto & \text{Cu}_A(-) \end{array}$$

is a covariant functor.

Theorem 4.2. *Let X be a one dimensional compact Hausdorff space and let A be a continuous field over X without K_1 obstructions. Consider the functors*

$$\begin{array}{ccc} \text{Cu}_A(-) : \mathcal{V}_X & \rightarrow & \text{Cu} & \text{and} & \Gamma(-, F_{\text{Cu}_A(-)}) : \mathcal{V}_X & \rightarrow & \text{Cu} \\ V & \mapsto & \text{Cu}(A(V)) & & V & \mapsto & \Gamma(V, F_{\text{Cu}_A(V)}) \end{array}$$

Then, $\text{Cu}_A(-)$ and $\Gamma(-, F_{\text{Cu}_A(-)})$ are isomorphic sheaves.

Proof. That $\text{Cu}_A(-)$ is a sheaf follows from Corollary 2.7. Let $(h_V)_{V \in \mathcal{V}_X}$ be the collection of isomorphisms $h_V : \text{Cu}(A(V)) \rightarrow \Gamma(V, F_{\text{Cu}_A(V)})$ described in Theorem 3.12. Since, whenever

$V \subset U$, the following diagram

$$\begin{array}{ccc} \mathrm{Cu}(A(V)) & \longrightarrow & \Gamma(V, F_{\mathrm{Cu}_A(V)}) \\ \uparrow (\mathrm{Cu}_A(\cdot))_V^U & & \uparrow (\Gamma(\cdot, F_{\mathrm{Cu}_A(\cdot)}))_V^U \\ \mathrm{Cu}(A(U)) & \longrightarrow & \Gamma(U, F_{\mathrm{Cu}_A(U)}) \end{array}$$

clearly commutes, $(h_V)_{V \in \mathcal{V}_X}$ defines an isomorphism of sheafs $h: \mathrm{Cu}_A(\cdot) \rightarrow \Gamma(\cdot, F_{\mathrm{Cu}_A(\cdot)})$. \square

In order to relate the Cuntz semigroup $\mathrm{Cu}(A)$ and the sheaf $\mathrm{Cu}_A(\cdot)$, we now show that there exists an action of $\mathrm{Cu}(C(X))$ on $\mathrm{Cu}(A)$ when A is a $C(X)$ -algebra, which is naturally induced from the $C(X)$ -module structure on A .

Definition 4.3. Let S, T, R be semigroups in Cu . A \ll -bimorphism is a map $\varphi: S \times T \rightarrow R$ such that the map $\varphi(s, \cdot): T \rightarrow R$, $s \in S$ (respectively, $\varphi(\cdot, t): S \rightarrow R$, $t \in T$), preserves order, addition, suprema of increasing sequences, and moreover $\varphi(s', t') \ll \varphi(s, t)$ whenever $s' \ll s$ in S and $t' \ll t$ in T .

Remark 4.4. We remark that if A is a C^* -algebra and a, b are commuting elements, then for $\epsilon > 0$ we have $(a - \epsilon)_+(b - \epsilon)_+ \preceq (ab - \epsilon^2)_+$. Indeed, since the C^* -subalgebra generated by a and b is commutative, Cuntz comparison is given by the support of the given elements, viewed as continuous functions on the spectrum of the algebra. It is then a simple matter to check that $\mathrm{supp}((a - \epsilon)_+(b - \epsilon)_+) \subseteq \mathrm{supp}((ab - \epsilon^2)_+)$.

Proposition 4.5. Let A and B be stable and nuclear C^* -algebras. Then, the natural bilinear map $A \times B \rightarrow A \otimes B$ given by $(a, b) \mapsto a \otimes b$ induces a \ll -bimorphism

$$\begin{array}{ccc} \mathrm{Cu}(A) \times \mathrm{Cu}(B) & \rightarrow & \mathrm{Cu}(A \otimes B) \\ ([a], [b]) & \mapsto & [a \otimes b] \end{array}$$

Proof. Since A is stable, we may think of $\mathrm{Cu}(A)$ as equivalence classes of positive elements from A . We also have an isomorphism $\Theta: M_2(A) \rightarrow A$ given by isometries w_1, w_2 in $\mathcal{M}(A)$ with orthogonal ranges, so that $\Theta(a_{ij}) = \sum_{i,j} w_i a_{ij} w_j^*$. Thus, in the Cuntz semigroup, $[a] + [b] = [\Theta(\begin{smallmatrix} a & 0 \\ 0 & b \end{smallmatrix})]$.

The map $\mathrm{Cu}(A) \times \mathrm{Cu}(B) \rightarrow \mathrm{Cu}(A \otimes B)$ given by $([a], [b]) \mapsto [a \otimes b]$ is well defined and order-preserving in each argument, by virtue of [27, Lemma 4.2]. Let $a, a' \in A_+$, $b \in B_+$. As

$$[(w_1 a w_1^* + w_2 a' w_2^*) \otimes b] = [w_1 a w_1^* \otimes b] + [w_2 a' w_2^* \otimes b] = [a \otimes b] + [a' \otimes b],$$

we see that it is additive in the first entry (and analogously in the second entry).

Next, observe that if $\|a\|, \|b\| \leq 1$, $\epsilon > 0$,

$$\|a \otimes b - (a - \epsilon)_+ \otimes (b - \epsilon)_+\| \leq \|a \otimes b - (a - \epsilon)_+ \otimes b\| + \|(a - \epsilon)_+ \otimes b - (a - \epsilon)_+ \otimes (b - \epsilon)_+\| \leq \epsilon \|b\| + \|(a - \epsilon)_+\| \epsilon \leq 2\epsilon,$$

and this implies $[a \otimes b] = \sup([(a - \epsilon)_+ \otimes (b - \epsilon)_+])$. If now $[a] = \sup_n [a_n]$ for an increasing sequence $([a_n])$, then for any $[b]$ we have $[a_n \otimes b] \leq [a \otimes b]$. Given $\epsilon > 0$, find n with $[(a - \epsilon)_+] \leq [a_n]$, hence $[(a - \epsilon)_+ \otimes (b - \epsilon)_+] \leq [a_n \otimes b] \leq \sup [a_n \otimes b]$. Taking supremum when ϵ goes to zero we obtain $[a \otimes b] = \sup [a_n \otimes b]$.

Finally, assume that $[a'] \ll [a]$ in $\text{Cu}(A)$, and $[b'] \ll [b]$ in $\text{Cu}(B)$. Find $\epsilon > 0$ such that $[a'] \leq [(a - \epsilon)_+]$ and $[b'] \leq [(b - \epsilon)_+]$. Then $[a' \otimes b'] \leq [(a - \epsilon)_+ \otimes (b - \epsilon)_+]$.

Note that $(a - \epsilon)_+ \otimes (b - \epsilon)_+ \in A \otimes B \subseteq \mathcal{M}(A) \otimes \mathcal{M}(B)$ and, viewed in the tensor product of the multiplier algebras, we have $(a - \epsilon)_+ \otimes (b - \epsilon)_+ = ((a - \epsilon)_+ \otimes 1)(1 \otimes (b - \epsilon)_+)$. Since $\mathcal{M}(A) \rightarrow \mathcal{M}(A) \otimes \mathcal{M}(B)$, $c \mapsto c \otimes 1$ is a $*$ -homomorphism, it induces a semigroup homomorphism $\text{Cu}(\mathcal{M}(A)) \rightarrow \text{Cu}(\mathcal{M}(A) \otimes \mathcal{M}(B))$ in the category Cu and, in particular, since $[(a - \epsilon)_+] \ll [a]$ in $\text{Cu}(\mathcal{M}(A))$, it follows that $[(a - \epsilon)_+ \otimes 1] \ll [a \otimes 1]$ in $\text{Cu}(\mathcal{M}(A) \otimes \mathcal{M}(B))$. Likewise, $[1 \otimes (b - \epsilon)_+] \ll [1 \otimes b]$, hence we may find $\epsilon' > 0$ such that $[(a - \epsilon)_+ \otimes 1] \leq [(a \otimes 1 - \epsilon')_+]$ and $[1 \otimes (b - \epsilon)_+] \leq [(1 \otimes b - \epsilon')_+]$. Since the elements $(a - \epsilon)_+ \otimes 1$, $(a \otimes 1 - \epsilon')_+$, $1 \otimes (b - \epsilon)_+$ and $(1 \otimes b - \epsilon')_+$ all commute (and using Remark 4.4), it follows that

$$\begin{aligned} [(a - \epsilon)_+ \otimes (b - \epsilon)_+] &= [((a - \epsilon)_+ \otimes 1)(1 \otimes (b - \epsilon)_+)] \\ &\leq [(a \otimes 1 - \epsilon')_+(1 \otimes b - \epsilon')_+] \\ &\leq [(a \otimes b - \epsilon'^2)_+] \ll [a \otimes b], \end{aligned}$$

whence $[a' \otimes b'] \ll [a \otimes b]$. □

Corollary 4.6. *Let X be a compact Hausdorff space, and let A be a stable $C(X)$ -algebra (with structure map θ). Then the natural map $C(X) \times A \rightarrow A$, given by $(f, a) \rightarrow \theta(f)a$ induces a \ll -bimorphism*

$$\gamma_A: \text{Cu}(C(X)) \times \text{Cu}(A) \rightarrow \text{Cu}(A)$$

such that maps $([f], [a])$ to $[\theta(f)a]$, for $f \in C(X)_+$ and $a \in A_+$.

Proof. Since $\text{Cu}(C(X)) = \text{Cu}(C(X) \otimes \mathcal{K})$, Proposition 4.5 tells us that the map

$$\begin{array}{ccc} \text{Cu}(C(X) \otimes \mathcal{K}) \times \text{Cu}(A) & \rightarrow & \text{Cu}(C(X) \otimes \mathcal{K} \otimes A) \\ ([f], [a]) & \mapsto & [f \otimes a] \end{array}$$

is a \ll -bimorphism. Now the result follows after composing this map with the isomorphism $\text{Cu}(C(X) \otimes \mathcal{K} \otimes A) \cong \text{Cu}(C(X) \otimes A)$, followed by the map $\text{Cu}(\theta): \text{Cu}(C(X) \otimes A) \rightarrow \text{Cu}(A)$. □

In what follows, we shall refer to the \ll -bimorphism γ_A above as the action of $\text{Cu}(C(X))$ on $\text{Cu}(A)$. If A and B are $C(X)$ -algebras, we will say then that a morphism $\varphi: \text{Cu}(A) \rightarrow \text{Cu}(B)$ preserves the action provided $\varphi(\gamma_A(x, y)) = \gamma_B(x, \varphi(y))$. Notice that this is always the case if φ is induced by a $*$ -homomorphism of $C(X)$ -algebras. We will write γ instead of γ_A , and we will moreover use the notation xy for $\gamma(x, y)$.

As noticed above, $\mathbb{V}_A(-)$ defines a continuous presheaf, and we show below that it becomes a sheaf when A does not have K_1 obstructions. For this we need a lemma (see [3]).

Lemma 4.7. *Let X be a one dimensional compact Hausdorff space and let $Y, Z \subseteq X$ be closed subsets of X . Let A be a continuous field over X without K_1 obstructions, and denote by $\pi_Y^Z: A(Y) \rightarrow A(Y \cap Z)$ and $\pi_Z^Y: A(Z) \rightarrow A(Y \cap Z)$ the quotient maps (given by restriction). Then, the map*

$$\beta: V(A(Y) \oplus_{A(Y \cap Z)} A(Z)) \rightarrow V(A(Y)) \oplus_{V(A(Y \cap Z))} V(A(Z))$$

defined by $\beta([(a, b)]) = ([a], [b])$ is an isomorphism.

Proof. We know from Corollary 2.7 that Cu_A is a sheaf in this case. Thus the map

$$\text{Cu}(A(Y) \oplus_{A(Y \cap Z)} A(Z)) \rightarrow \text{Cu}(A(Y)) \oplus_{\text{Cu}(A(Y \cap Z))} \text{Cu}(A(Z)),$$

given by $[(a, b)] \mapsto ([a], [b])$, is an isomorphism in Cu , whence it maps compact elements to compact elements. Since $A(Y) \oplus_{A(Y \cap Z)} A(Z)$ is isomorphic to $A(Y \cup Z)$ and this algebra is stably finite (because all of its fibers have stable rank one), we have that the compact elements of $\text{Cu}(A(Y) \oplus_{A(Y \cap Z)} A(Z))$ can be identified with $V(A(Y) \oplus_{A(Y \cap Z)} A(Z))$.

Using this identification, we have that $[(a, b)]$ in $\text{Cu}(A(Y) \oplus_{A(Y \cap Z)} A(Z))$ is compact if and only if $[a]$ and $[b]$ are compact. On the other hand, if $[a]$ and $[b]$ are compact (in $\text{Cu}(A(Y))$ and $\text{Cu}(A(Z))$ respectively) and $[a] = [b]$ in $\text{Cu}(A(Y \cap Z))$, then the pair $([a], [b])$ belongs to $V(A(Y)) \oplus_{V(A(Y \cap Z))} V(A(Z))$, and every element of this pullback is obtained in this manner. The conclusion now follows easily. \square

Proposition 4.8. *Let X be a one dimensional compact Hausdorff space and let A be a continuous field over X without K_1 obstructions. Then,*

$$\begin{array}{ccc} \mathbb{V}_A(-) : \mathcal{V}_X & \rightarrow & \text{Sg} \\ & U \mapsto & V(A(U)) \end{array}$$

is a sheaf and the natural transformation $\mathbb{V}_A(-) \rightarrow \Gamma(-, F_{V(A(\cdot))})$ is an isomorphism of sheaves.

Proof. Note that $\mathbb{V}_A(-)$ is a sheaf thanks to Lemma 4.7. On the other hand, the fact that $\mathbb{V}_A(-)$ is isomorphic to the sheaf of continuous sections $\Gamma(-, F_{V(A(\cdot))})$ follows from Theorem 2.2 in [30]. \square

Theorem 4.9. *Let X be a compact Hausdorff space and let A and B be $C(X)$ -algebras such that all fibers have stable rank one. Consider the following conditions:*

- (i) $\text{Cu}(A) \cong \text{Cu}(B)$ preserving the action of $\text{Cu}(C(X))$,
- (ii) $\text{Cu}_A(-) \cong \text{Cu}_B(-)$,
- (iii) $\mathbb{V}_A(-) \cong \mathbb{V}_B(-)$.

Then (i) \implies (ii) \implies (iii). If X is one dimensional, then also (ii) \implies (i). If, furthermore, A and B are continuous fields without K_1 obstructions such that for all $x \in X$ the fibers A_x, B_x have real rank zero, then (iii) \implies (ii) and so all three conditions are equivalent.

Proof. We may assume that both A and B are stable.

(i) \implies (ii): Let $\varphi : \text{Cu}(A) \rightarrow \text{Cu}(B)$ be an isomorphism such that $\varphi(xy) = x\varphi(y)$, for any $x \in \text{Cu}(C(X))$ and $y \in \text{Cu}(A)$. We need to verify that $\varphi(\text{Cu}(C_0(X \setminus V)A)) \subseteq \text{Cu}(C_0(X \setminus V)B)$, whenever V is a closed subset of X . This will entail that φ induces a semigroup map $\varphi_V : \text{Cu}(A(V)) \rightarrow \text{Cu}(B(V))$, which is an isomorphism as φ is.

Let $[fa] \in \text{Cu}(C_0(X \setminus V)A)$, for $f \in C_0(X \setminus V)_+$ and $a \in A_+$. Then, if $\varphi([a]) = [b]$ for some $b \in B_+$, we have that $\varphi([fa]) = [f]\varphi([a]) = [f][b] = [fb]$, and $fb \in C_0(X \setminus V)B$. Thus $\varphi(\text{Cu}(C_0(X \setminus V)A)) \subseteq \varphi(C_0(X \setminus V)B)$. The rest of the argument is routine.

(ii) \implies (iii): Note that, as all fibers have stable rank one, $\text{Cu}(A(U))$ (respectively, $\text{Cu}(B(U))$) is a stably finite algebra for each closed subset U . In this case, $V(A(U))$ can be identified with the subset of compact elements of $\text{Cu}(A(U))$. Therefore, the given isomorphism $\text{Cu}_A(U) \cong \text{Cu}_B(U)$ maps $\mathbb{V}_A(U) = V(A(U))$ injectively onto $\mathbb{V}_B(U) = V(B(U))$.

Now assume that X is one dimensional, and let us prove that (ii) \implies (i): The isomorphism of sheaves gives, in particular, an isomorphism $\varphi : \text{Cu}(A) \rightarrow \text{Cu}(B)$. We need to

verify that φ respects the action of $\text{Cu}(C(X))$. By [21], $\text{Cu}(C(X)) \cong \text{Lsc}(X, \overline{\mathbb{N}})$. In this case, any $f \in \text{Lsc}(X, \overline{\mathbb{N}})$ may be written as:

$$f = \sum_{i=1}^{\infty} \mathbb{1}_{U_i} \text{ where } U_i = f^{-1}((i, \infty]).$$

Thus, in order to check that $\varphi(f[a]) = f\varphi([a])$, it is enough to verify it when $f = \mathbb{1}_U$, where U is an open set of X .

Notice that $\mathbb{1}_U[a] = [ga]$ where $g \in C(X)_+$ has $\text{supp}(g) = U$. Given $[a] \in \text{Cu}(A)$ we denote by $\text{supp}([a]) = \{x \in X \mid \pi_x([a]) \neq 0\}$. Observe that $\text{supp}\varphi([a]) = \text{supp}([a])$, and that $\text{supp}(\mathbb{1}_U\varphi([a])) = U \cap \text{supp}([a])$. Let $K \subseteq \text{supp}(\mathbb{1}_U\varphi([a])) = \text{supp}(\varphi(\mathbb{1}_U[a]))$ be a closed set. Then $\pi_K(\mathbb{1}_U\varphi([a])) = \pi_K(\varphi(\mathbb{1}_U[a]))$, where $\pi_K: A \rightarrow A(K)$ is the quotient map. Indeed, it follows from the commutative diagram

$$\begin{array}{ccc} \text{Cu}(A) & \xrightarrow{\varphi} & \text{Cu}(B) \\ \pi_K \downarrow & & \downarrow \pi_K \\ \text{Cu}(A(K)) & \xrightarrow{\varphi_K} & \text{Cu}(B(K)) \end{array},$$

that $\pi_K(\mathbb{1}_U[a]) = \pi_K([ga]) = \pi_K[a]$, since g becomes invertible in $A(K)$. Therefore,

$$\pi_K(\varphi(\mathbb{1}_U[a])) = \pi_K(\varphi([ga])) = \varphi_K\pi_K([ga]) = \varphi_K\pi_K([a]).$$

On the other hand, $\pi_K(\mathbb{1}_U\varphi([a])) = \pi_K\varphi([a]) = \varphi_K\pi_K([a])$.

Now write $[a] = \sup[a_n]$, where $([a_n])$ is a rapidly increasing sequence in $\text{Cu}(A)$, and $\mathbb{1}_U = \sup \mathbb{1}_{V_n}$, where (V_n) is a rapidly increasing sequence of open sets. Then $(\mathbb{1}_{V_n}[a_n])$ is a rapidly increasing sequence with $\mathbb{1}_U[a] = \sup \mathbb{1}_{V_n}[a_n]$ and $\mathbb{1}_U\varphi([a]) = \sup \mathbb{1}_{V_n}\varphi([a_n])$. By [2, Lemma 2.5] choose, for each n , a compact set K_n such that

$$\text{supp}(\mathbb{1}_{V_n}[a_n]) \subseteq K_n \subseteq \text{supp}(\mathbb{1}_{V_{n+1}}[a_{n+1}]).$$

Then $K_n \subseteq V_{n+1} \cap \text{supp}([a_{n+1}]) \subseteq V_{n+1}$.

By the above, $\pi_{K_n}(\mathbb{1}_{V_{n+1}}\varphi([a_{n+1}])) = \pi_{K_n}\varphi(\mathbb{1}_{V_{n+1}}[a_{n+1}])$, and thus:

$$\pi_{K_n}(\mathbb{1}_{V_n}\varphi([a_n])) \leq \pi_{K_n}(\mathbb{1}_{V_{n+1}}\varphi([a_{n+1}])) = \pi_{K_n}(\varphi(\mathbb{1}_{V_{n+1}}[a_{n+1}])) \leq \pi_{K_n}(\varphi(\mathbb{1}_U[a])),$$

and

$$\pi_{K_n}(\varphi(\mathbb{1}_{V_n}[a_n])) \leq \pi_{K_n}(\varphi(\mathbb{1}_{V_{n+1}}[a_{n+1}])) = \pi_{K_n}(\mathbb{1}_{V_{n+1}}\varphi([a_{n+1}])) \leq \pi_{K_n}(\mathbb{1}_U\varphi([a])).$$

Since $\text{supp}(\mathbb{1}_{V_k}[a_k]) = \text{supp}(\varphi(\mathbb{1}_{V_k}[a_k])) = \text{supp}(\mathbb{1}_{V_k}\varphi([a_k]))$, we may apply Lemma 2.4 in [2] to obtain that $\mathbb{1}_{V_n}\varphi([a_n]) \leq \varphi(\mathbb{1}_U[a])$ and $\varphi(\mathbb{1}_{V_n}[a_n]) \leq \mathbb{1}_U\varphi([a])$. Taking suprema in both inequalities we obtain $\mathbb{1}_U\varphi([a]) = \varphi(\mathbb{1}_U[a])$.

(iii) \implies (ii): We assume now that both A and B are continuous fields without K_1 obstructions such that A_x and B_x have real rank zero for all x . Let $\varphi: \mathbb{V}_A(-) \rightarrow \mathbb{V}_B(-)$ be a sheaf isomorphism. This induces a semigroup isomorphism $\varphi_x: V(A_x) \rightarrow V(B_x)$ for each $x \in X$. As $A(U)$ is a stably finite algebra for any closed subset U of X , we will identify $V(A(U))$ with its image in $\text{Cu}(A(U))$ whenever convenient.

Since A_x has real rank zero, $V(A_x)$ forms a dense subset of $\text{Cu}(A_x)$ so we can uniquely define an isomorphism $\text{Cu}(A_x) \rightarrow \text{Cu}(B_x)$ in Cu which we will still denote by φ_x . This map is defined by $\varphi_x(z) = \sup_n \varphi_x(z_n)$ where $z = \sup z_n$ and $z_n \in V(A_x)$ for all $n \geq 0$ (see, e.g.

[1], [8] for further details). Let us prove that the induced bijective map $\tilde{\varphi}: F_{\text{Cu}(A)} \rightarrow F_{\text{Cu}(B)}$ is continuous, and hence an homeomorphism. This will define an isomorphism of sheaves $\Gamma(-, F_{\text{Cu}_A(-)}) \cong \Gamma(-, F_{\text{Cu}_B(-)})$ and then, using Theorem 4.2, it follows that $\text{Cu}_A(-)$ and $\text{Cu}_B(-)$ are isomorphic.

Denote by $\pi_A: F_{\text{Cu}(A)} \rightarrow X$ and $\pi_B: F_{\text{Cu}(B)} \rightarrow X$ the natural maps. Let U be an open set of X and $s \in \text{Cu}(B)$. We are to show that $\tilde{\varphi}^{-1}(U_s^{\gg})$ is open in $F_{\text{Cu}(A)}$. Let $z \in \tilde{\varphi}^{-1}(U_s^{\gg})$, and put $x = \pi_A(z)$, so that $z \in \text{Cu}(A_x)$ for some $x \in U$. Since $\tilde{\varphi}(z) = \varphi_x(z) \in U_s^{\gg}$, there exists $s'' \gg s$ such that $\hat{s}''(x) \ll \varphi_x(z)$. Choose s' such that $s \ll s' \ll s''$.

As $\hat{s}''(x) \ll \varphi_x(z)$ there exists $z' \ll z' \in V(A_x)$ such that $\hat{s}''(x) \ll \varphi_x(z')$. Now we can find a closed subset W' whose interior contains x , and an element $v \in V(A(W'))$ such that $\pi_x(v) = z'$. Note that $\pi_x \varphi_{W'}(v) = \varphi_x(z')$. Also, since $\hat{s}''(x) \ll \hat{s}''(x) \ll \varphi_x(z') = \widehat{\varphi_{W'}(v)}(x)$, we may use Lemma 2.2 to find $W \subseteq W'$ such that $x \in \overset{\circ}{W}$ and

$$\hat{s}''(y) \ll \widehat{\varphi_{W'}(v)}(y) \text{ for all } y \in W.$$

Let $t \in \text{Cu}(A)$ be such that $\pi_W(t) = \pi_W^{\overset{\circ}{W}}(v)$. We now claim that $\overset{\circ}{W}_t^{\gg} \subseteq \tilde{\varphi}^{-1}(U_s^{\gg})$. Let $w \in \overset{\circ}{W}_t^{\gg}$, and put $y = \pi_A(w) \in W$. There exists $t' \gg t$ such that $\hat{t}'(y) \ll w$, whence, applying $\tilde{\varphi}$ it follows that

$$\tilde{\varphi}(w) \gg \tilde{\varphi}(\hat{t}'(y)) \gg \tilde{\varphi}(\hat{t}(y)) = \tilde{\varphi}(\pi_W^{\overset{\circ}{W}}(v)(y)) = \tilde{\varphi}(\pi_y(v)) = \pi_y(\varphi_W(v)) = \widehat{\varphi_{W'}(v)}(y) \gg \hat{s}''(y),$$

and this shows that $w \in \tilde{\varphi}^{-1}(U_s^{\gg})$. □

Remark 4.10. We remark that the implication (ii) \implies (i) in Theorem 4.9 above holds whenever $\text{Cu}(C(X)) \cong \text{Lsc}(X, \overline{\mathbb{N}})$. This is the case for spaces more general than being just one dimensional, see [21].

Remark 4.11. Our Theorem 4.9 above allows us to rephrase the classification result obtained in [14], by using a single invariant. Namely, let A, B be separable unital continuous fields of AF -algebras over $[0, 1]$, and let $\tilde{\varphi}: \text{Cu}(A) \rightarrow \text{Cu}(B)$ be an isomorphism that preserves the action by $\text{Lsc}([0, 1], \overline{\mathbb{N}})$ and such that $\tilde{\varphi}([1_A]) = [1_B]$. Then $\tilde{\varphi}$ lifts to an isomorphism $\phi: A \rightarrow B$ of continuous fields of C^* -algebras.

ACKNOWLEDGEMENTS

This work has been partially supported by a DGI MICIIN grant (Spain) through Project MTM2011-28992-C02-01, and by the Comissionat per Universitats i Recerca de la Generalitat de Catalunya.

REFERENCES

- [1] R. Antoine, J. Bosa, F. Perera, *Completions of monoids with applications to the Cuntz Semigroup*, Internat. J. Math., **22**(5), 837–861, 2011.
- [2] R. Antoine, M. Dadarlat, F. Perera, L. Santiago, *Recovering the Elliott invariant from the Cuntz semigroup*, Trans. Amer. Math. Soc. (to appear).
- [3] R. Antoine, F. Perera, L. Santiago, *Pullbacks, $C(X)$ -algebras, and their Cuntz Semigroup* J. Funct. Anal. **260**(10), 2844–2880, 2011.
- [4] P. Ara, F. Perera, A. S. Toms, *K-theory for operator algebras. Classification of C^* -algebras* (in “Aspects of Operator Algebras and Applications”). Contemp. Math. **534**, 1–71, Amer. Math. Soc. 2011.

- [5] B. Blackadar, L. Robert, A. P. Tikuisis, A. S. Toms, W. Winter, *An algebraic approach to the radius of comparison*, Trans. Amer. Math. Soc., **364**, 3657–3674, 2012.
- [6] E. Blanchard, *Déformations de C^* -algèbres de Hopf*, Bull. Soc. Math. France, **124**, 141–215, 1996.
- [7] N. P. Brown, F. Perera, and A. S. Toms, *The Cuntz semigroup, the Elliott conjecture, and dimension functions on C^* -algebras*, J. Reine Angew. Math., **621**, 191–211, 2008.
- [8] K. T. Coward, G. A. Elliott and C. Ivanescu, *The Cuntz Semigroup as an invariant for C^* -algebras*, J. Reine Angew. Math., **623**, 161–193, 2008.
- [9] A. Ciuperca, G. A. Elliott, *A remark on invariants for C^* -algebras of stable rank one*, Int. Math. Res. Not. IMRN, **5**, Art. ID rnm 158, 33 pp, 2008.
- [10] A. Ciuperca, G. A. Elliott, L. Santiago, *On inductive limits of type-I C^* -algebras with one-dimensional spectrum*, Int. Math. Res. Not. IMRN, **11**, 2577–2615, 2011.
- [11] A. Ciuperca, L. Robert, L. Santiago, *The Cuntz semigroup of ideals and quotients and a generalized Kasparov stabilization theorem*, J. Operator Theory, **64**, 155–169, 2010.
- [12] M. Dadarlat, *Continuous fields of C^* -algebras over finite dimensional spaces*, Adv. Math., **222**, 1850–1881, 2009.
- [13] M. Dadarlat, G. A. Elliott, *One-Parameter Continuous Fields of Kirchberg Algebras*, Comm. Math. Phys. **274**, 795–819, 2007.
- [14] M. Dadarlat, G. A. Elliott, Z. Niu, *One-Parameter Continuous Fields of Kirchberg Algebras II*, Canad. J. Math. **63** (3), 500–532, 2011.
- [15] J. Dixmier, *C^* -algebras*, North-Holland Mathematical Library, **15**, North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977.
- [16] E. Kirchberg, M. Rørdam, *Infinite non-simple C^* -algebras: absorbing the Cuntz algebras \mathcal{O}_∞* , Adv. Math., **167** (2), 198–264, 2002.
- [17] M. Nilsen, *C^* -bundles and $C_0(X)$ -algebras*, Indiana Univ. Math. J., **45** (2), 463–477, 1996.
- [18] C. Pasnicu, *Real rank zero and continuous fields of C^* -algebras*, Bull. Math. Soc. Sci. Math. Roumanie (N.S.), **48(96)** (3), 319–325, 2005.
- [19] C. Pasnicu, *The ideal property, the projection property, continuous fields and crossed products*, J. Math. Anal. Appl. **323** (2), 1213–1224, 2006.
- [20] A. Pears, *Dimension Theory of General Spaces*, Cambridge University Press, 1975.
- [21] L. Robert, *The Cuntz semigroup of some spaces of dimension at most 2*, C. R. Math. Acad. Sci. Soc. R. Can., to appear.
- [22] L. Robert, *The cone of functionals on the Cuntz semigroup*, Math. Scand., to appear.
- [23] L. Robert, *Classification of inductive limits of 1-dimensional NCCW complexes*, Adv. in Math., **231** (5), 2802–2836, 2012.
- [24] L. Robert, A. P. Tikuisis, *Hilbert C^* -modules over a commutative C^* -algebra*, Proc. Lond. Math. Soc. **102** (2), 229–256, 2011.
- [25] M. Rørdam, *On the structure of simple C^* -algebras tensored with a UHF-algebra II*, J. Funct. Anal., **107** (2), 255–269, 1992.
- [26] M. Rørdam, *A simple C^* -algebra with a finite and an infinite projection*, Acta Math., **191** (1), 109–142, 2003.
- [27] M. Rørdam, *The stable and the real rank of \mathcal{Z} -absorbing C^* -algebras*, Internat. J. Math., **15** (10), 1065–1084, 2004.
- [28] A. Tikuisis, *The Cuntz semigroup of continuous functions into certain simple C^* -algebras*, Internat. J. Math., **22** (8), 1051–1087, 2011.
- [29] A. S. Toms, *On the classification problem for nuclear C^* -algebras*, Ann. of Math. (2), **167** (3), 1029–1044, 2008.
- [30] R. O. Jr. Wells, *Differential analysis on complex manifolds*, Graduate Texts in Mathematics **65**, Springer-Verlag, 1980.
- [31] N. E. Wegge-Olsen, *K -theory and C^* -algebras*, Oxford Science Publications. The Clarendon Press, Oxford University Press, 1993.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 BELLATERRA, BARCELONA, SPAIN

E-mail address: ramon@mat.uab.cat, jbosa@mat.uab.cat, perera@mat.uab.cat