



Discrete Laguerre–Sobolev expansions: A Cohen type inequality[☆]

A. Peña^{*}, M.L. Rezola

Departamento de Matemáticas and IUMA, Universidad de Zaragoza, 50009 Zaragoza, Spain

ARTICLE INFO

Article history:

Received 11 March 2011

Available online 16 June 2011

Submitted by D. Khavinson

Keywords:

Laguerre polynomials

Laguerre–Sobolev type polynomials

Cohen type inequality

Bessel functions

ABSTRACT

C. Markett proved a Cohen type inequality for the classical Laguerre expansions in the appropriate weighted L^p spaces. In this paper, we get a Cohen type inequality for the Fourier expansions in terms of discrete Laguerre–Sobolev orthonormal polynomials with an arbitrary (finite) number of mass points. So, we extend the result due to B.Xh. Fejzullahu and F. Marcellán.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction and notations

Littlewood conjectured in 1948 that for any trigonometric polynomial $F_N(x) = \sum_{k=1}^N a_k e^{in_k x}$ where $0 < n_1 < n_2 < \dots < n_N$, $N \geq 2$, and $|a_k| \geq 1$ for $1 \leq k \leq N$, there holds the estimate from below

$$\int_0^{2\pi} |F_N(x)| dx \geq C \log N$$

where C is an absolute constant (see [6]).

Cohen's inequality [3] was the first result on the way to the solution of this conjecture. Later, inequalities of this type have been established in various other contexts, e.g., on compact group (see [5]).

In [12] Markett proved such inequalities for classical orthogonal polynomial expansions in the appropriate weighted L^p spaces, here in terms of the highest coefficient. The main purpose of this paper is to extend these results to discrete Laguerre–Sobolev expansions. More precisely, we obtain such inequalities, in the appropriate weighted L^p spaces, for Fourier expansions in terms of orthonormal polynomials with respect to an inner product of the form

$$\langle p, q \rangle_s = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty p(x)q(x)x^\alpha e^{-x} dx + \sum_{j=0}^N M_j p^{(j)}(0)q^{(j)}(0), \quad (1)$$

where $\alpha > -1$ and $M_j \geq 0$, $j = 0, \dots, N$. Such inner products are called of discrete Sobolev type.

Recently in [4], the authors Fejzullahu and Marcellán obtained Cohen type inequalities for orthonormal expansions with respect to the above inner product in the case $N = 1$, i.e. at most two masses in the discrete part. In this particular

[☆] Both authors partially supported by MICINN of Spain under Grant MTM2009-12740-C03-03, FEDER funds (EU), and the DGA project E-64 (Spain).

^{*} Corresponding author. Fax: +34 976761338.

E-mail address: anap@unizar.es (A. Peña).

case, the authors benefit from the fact that there are explicit formulas for the connection coefficients which appear in the representation of discrete Laguerre–Sobolev type polynomials in terms of three standard Laguerre polynomials (see [9]). For a general discrete Laguerre–Sobolev inner product, we only know that these coefficients are a nontrivial solution of a system of $N + 1$ equations on $N + 2$ unknowns (see [8]). If the system is solved, we get an intricate expression with which it is difficult to work. Our contribution in this paper is that we can assure that there exists limit of these connection coefficients and this is enough for our purpose.

Let $\{L_n^\alpha(x)\}_{n \geq 0}$ be the sequence of Laguerre polynomials, orthogonal on $[0, \infty)$ with respect to the probability measure $d\mu(x) = \frac{1}{\Gamma(\alpha+1)} x^\alpha e^{-x} dx$ where $\alpha > -1$ and normalized by $L_n^\alpha(0) = \binom{n+\alpha}{n}$. We denote the orthonormal Laguerre polynomial of degree n by

$$l_n^\alpha(x) = \frac{L_n^\alpha(x)}{\|L_n^\alpha\|}$$

where $\|L_n^\alpha\|^2 = \int_0^\infty L_n^\alpha(x)^2 d\mu(x)$.

Let $\{Q_n^\alpha\}_{n \geq 0}$ be the sequence of discrete Laguerre–Sobolev orthogonal polynomials with respect to the inner product (1) and such that $Q_n^\alpha(x)$ and $L_n^\alpha(x)$ have the same leading coefficient. We denote by

$$q_n^\alpha(x) = \langle Q_n^\alpha, Q_n^\alpha \rangle_S^{-1/2} Q_n^\alpha(x)$$

the orthonormal discrete Laguerre–Sobolev polynomials. From now on, for simplicity we write $Q_n(x) = Q_n^\alpha(x)$ and $q_n(x) = q_n^\alpha(x)$.

Laguerre expansions have been investigated mainly in the following two sets of weighted Lebesgue spaces, namely in the classical spaces [2,10]

$$L_p(x^\alpha dx) = \begin{cases} \{f; \int_0^\infty |f(x)e^{-x/2}|^p x^\alpha dx < \infty\}, & \text{if } 1 \leq p < \infty; \\ \{f; \text{ess sup}_{0 < x < \infty} |f(x)e^{-x/2}| < \infty\}, & \text{if } p = \infty, \end{cases}$$

for $\alpha > -1$ as well as in the spaces

$$L_p(x^{\alpha p/2} dx) = \begin{cases} \{f; \int_0^\infty |f(x)e^{-x/2} x^{\alpha/2}|^p dx < \infty\}, & \text{if } 1 \leq p < \infty; \\ \{f; \text{ess sup}_{0 < x < \infty} |f(x)e^{-x/2} x^{\alpha/2}| < \infty\}, & \text{if } p = \infty, \end{cases}$$

for $\alpha > -\frac{2}{p}$ if $1 \leq p < \infty$ and $\alpha \geq 0$ if $p = \infty$.

In order to unify the two results we are going to prove, we introduce an auxiliary parameter β which means either α or $\alpha p/2$.

We consider the class S_p^β , $1 \leq p \leq \infty$, defined as the space of measurable functions f defined on $[0, \infty)$, such that there exists $f^{(k)}(0)$ for $k = 0, \dots, N$ and if $1 \leq p < \infty$

$$\|f\|_{S_p^\beta}^p = \|f\|_{L_p(x^\beta dx)}^p + \sum_{j=0}^N M_j |f^{(j)}(0)|^p < \infty,$$

where

$$\|f\|_{L_p(x^\beta dx)}^p = \int_0^\infty |f(x)e^{-x/2}|^p x^\beta dx, \quad 1 \leq p < \infty,$$

and if $p = \infty$

$$\|f\|_{S_\infty^\beta} = \max\{\|f\|_{L_\infty(x^\beta dx)}, |f(0)|, \dots, |f^{(N)}(0)|\} < \infty,$$

where

$$\|f\|_{L_\infty(x^\beta dx)} = \begin{cases} \text{ess sup}_{0 < x < \infty} |f(x)e^{-x/2}|, & \text{if } \beta = \alpha; \\ \text{ess sup}_{0 < x < \infty} |f(x)e^{-x/2} x^{\alpha/2}|, & \text{if } \beta = \alpha p/2. \end{cases}$$

(If some $M_j = 0$ the corresponding derivative does not appear in the maximum.)

Let $f \in S_p^\beta$, $1 \leq p \leq \infty$, then the Fourier expansion in terms of orthonormal discrete Laguerre–Sobolev polynomials $\{q_n\}_{n \geq 0}$, is

$$\sum_{k=0}^\infty \hat{f}(k) q_k(x)$$

where $\hat{f}(k) = \langle f, q_k \rangle_S$.

In the following, $[S_p^\beta]$ denotes the space of all bounded linear operators T from the space S_p^β into itself, endowed with the usual operator norm,

$$\|T\|_{[S_p^\beta]} = \sup_{0 \neq f \in S_p^\beta} \frac{\|Tf\|_{S_p^\beta}}{\|f\|_{S_p^\beta}}.$$

Let $1 \leq p \leq \infty$. For a family of complex numbers $\{c_{k,n}\}_{k=0}^n$, $n \in \mathbb{N} \cup \{0\}$, with $|c_{n,n}| > 0$ we define the operators $T_n^{\alpha,S} : S_p^\beta \rightarrow S_p^\beta$ by

$$T_n^{\alpha,S}(f) = \sum_{k=0}^n c_{k,n} \hat{f}(k) q_k.$$

Let us denote $q_0 = \frac{4\alpha+4}{2\alpha+1}$ for $\beta = \alpha$ and $q_0 = 4$ for $\beta = p\alpha/2$, and let p_0 be the conjugate of q_0 , i.e. $1/p_0 + 1/q_0 = 1$. Now, we can state our main theorem, which extends the ones given in [12] and [4].

Theorem 1. *Let $1 \leq p \leq \infty$. There exists a positive constant C , independent of n , such that:
For $\alpha > -1/2$*

$$\|T_n^{\alpha,S}\|_{[S_p^\beta]} \geq C|c_{n,n}| \begin{cases} n^{\frac{2\alpha+2}{p} - \frac{2\alpha+3}{2}}, & \text{if } 1 \leq p < p_0; \\ (\log(n+1))^{\frac{2\alpha+1}{4\alpha+4}}, & \text{if } p = p_0, p = q_0; \\ n^{\frac{2\alpha+1}{2} - \frac{2\alpha+2}{p}}, & \text{if } q_0 < p \leq \infty. \end{cases}$$

For $\alpha > -2/p$ if $1 \leq p < \infty$ and $\alpha \geq 0$ if $p = \infty$

$$\|T_n^{\alpha,S}\|_{[S_{p_0}^{q_0/2}]} \geq C|c_{n,n}| \begin{cases} n^{\frac{2}{p} - \frac{3}{2}}, & \text{if } 1 \leq p < p_0; \\ (\log(n+1))^{\frac{1}{4}}, & \text{if } p = p_0, p = q_0; \\ n^{\frac{1}{2} - \frac{2}{p}}, & \text{if } q_0 < p \leq \infty. \end{cases}$$

This theorem will be proved in Section 3. In Section 2, we obtain some new results for discrete Laguerre–Sobolev polynomials, which we will use to establish Theorem 1. More concretely, we prove a technical lemma that will be used to deduce a Mehler–Heine type formula for Laguerre–Sobolev polynomials and a sharp estimation for their norm in the appropriate weighted L_p spaces.

In the sequel we use the following notation, $a_n \sim b_n$ means that there exist positive constants c_1 and c_2 , such that $c_1 a_n \leq b_n \leq c_2 a_n$ for n large enough, while $a_n \cong b_n$ means that the sequence $\frac{a_n}{b_n}$ converges to 1. Throughout the paper, the values of the constants may change from line to line.

2. Estimates for discrete Laguerre–Sobolev polynomials

Consider the standard Laguerre polynomials L_n^α and the Laguerre–Sobolev polynomials Q_n with the same leading coefficient.

Let us recall some properties of Laguerre polynomials for $\alpha > -1$ (see [14]). The evaluation at $x = 0$ of the polynomials L_n^α and its successive derivatives are given by

$$(L_n^\alpha)^{(k)}(0) = \frac{(-1)^k \Gamma(n + \alpha + 1)}{(n - k)! \Gamma(\alpha + k + 1)}, \quad k \in \mathbb{N} \cup \{0\},$$

and their L_2 -norm is

$$\|L_n^\alpha\|^2 = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty (L_n^\alpha(x))^2 x^\alpha e^{-x} dx = \frac{\Gamma(n + \alpha + 1)}{n! \Gamma(\alpha + 1)}. \tag{2}$$

As usual, we denote the derivatives of the n th kernels of Laguerre polynomials by

$$K_n^{(k,h)}(x, y) = \frac{\partial^{k+h}}{\partial x^k \partial y^h} K_n(x, y) = \sum_{i=0}^n \frac{(L_i^\alpha)^{(k)}(x) (L_i^\alpha)^{(h)}(y)}{\|L_i^\alpha\|^2}$$

with $k, h \in \mathbb{N} \cup \{0\}$ and the convention $K_n^{(0,0)}(x, y) = K_n(x, y)$.

In the next lemma, we obtain an asymptotic estimate for $Q_n^{(k)}(0)$, that will play an important role along this paper.

Lemma 1. *Let Q_n be the polynomials orthogonal with respect to the inner product (1). Then the following statements hold:*

(a)

$$\frac{Q_n^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} \cong \begin{cases} \frac{C_k}{n^{\alpha+2k+1}}, & \text{for } k \text{ such that } M_k > 0; \\ C_k, & \text{otherwise,} \end{cases}$$

where C_k is a nonzero constant independent of n .

(b)

$$\langle Q_n, Q_n \rangle_S \cong \|L_n^\alpha\|^2.$$

Proof. If all the masses in the inner product (1) are zero the result is trivial because $Q_n = L_n^\alpha$. We will prove the result by induction concerning the number of positive masses in the inner product (1).

We take the first mass which is positive, namely M_{j_1} ($j_1 \geq 0$), and consider the sequence of polynomials $\{Q_{n,1}\}_{n \geq 0}$ orthogonal with respect to the inner product

$$(p, q)_1 = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty p(x)q(x)x^\alpha e^{-x} dx + M_{j_1} p^{(j_1)}(0)q^{(j_1)}(0).$$

The Fourier expansion of the polynomial $Q_{n,1}$ in the orthogonal basis $\{L_n^\alpha\}_{n \geq 0}$ leads to

$$Q_{n,1}(x) = L_n^\alpha(x) - M_{j_1} Q_{n,1}^{(j_1)}(0) K_{n-1}^{(0,j_1)}(x, 0).$$

Therefore

$$Q_{n,1}(x) = L_n^\alpha(x) - \frac{M_{j_1} (L_n^\alpha)^{(j_1)}(0)}{1 + M_{j_1} K_{n-1}^{(j_1,j_1)}(0, 0)} K_{n-1}^{(0,j_1)}(x, 0), \tag{3}$$

and

$$\langle Q_{n,1}, Q_{n,1} \rangle_1 = \|L_n^\alpha\|^2 + M_{j_1} \frac{((L_n^\alpha)^{(j_1)}(0))^2}{1 + M_{j_1} K_{n-1}^{(j_1,j_1)}(0, 0)}. \tag{4}$$

These relationships are very well known in the literature of discrete Sobolev type orthogonal polynomials.

Taking derivatives k times in (3) and evaluating at $x = 0$, we obtain

$$\frac{Q_{n,1}^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} = 1 - \frac{M_{j_1} K_{n-1}^{(k,j_1)}(0, 0)}{1 + M_{j_1} K_{n-1}^{(j_1,j_1)}(0, 0)} \frac{(L_n^\alpha)^{(j_1)}(0)}{(L_n^\alpha)^{(k)}(0)}. \tag{5}$$

Applying the Stolz criterion (see, e.g. [7]), we have

$$\lim_n \frac{K_{n-1}^{(k,j_1)}(0, 0)}{n^{\alpha+k+j_1+1}} = \lim_n \frac{(L_{n-1}^\alpha)^{(k)}(0)(L_{n-1}^\alpha)^{(j_1)}(0)}{\|L_{n-1}^\alpha\|^2(\alpha + k + j_1 + 1)n^{\alpha+k+j_1}} \neq 0, \tag{6}$$

and therefore

$$\frac{K_{n-1}^{(k,j_1)}(0, 0)}{K_{n-1}^{(j_1,j_1)}(0, 0)} \frac{(L_n^\alpha)^{(j_1)}(0)}{(L_n^\alpha)^{(k)}(0)} \cong \frac{(\alpha + 2j_1 + 1)}{(\alpha + k + j_1 + 1)} \frac{(L_{n-1}^\alpha)^{(k)}(0)}{(L_{n-1}^\alpha)^{(j_1)}(0)} \frac{(L_n^\alpha)^{(j_1)}(0)}{(L_n^\alpha)^{(k)}(0)} \cong \frac{\alpha + 2j_1 + 1}{\alpha + k + j_1 + 1}. \tag{7}$$

Thus, from (5), (6) and (7), we have

$$\frac{Q_{n,1}^{(j_1)}(0)}{(L_n^\alpha)^{(j_1)}(0)} = \frac{1}{1 + M_{j_1} K_{n-1}^{(j_1,j_1)}(0, 0)} \cong \frac{C_{j_1}}{n^{\alpha+2j_1+1}}$$

and for $k \neq j_1$

$$\frac{Q_{n,1}^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} \cong 1 - \frac{\alpha + 2j_1 + 1}{\alpha + k + j_1 + 1} \neq 0.$$

So, we achieve (a) for $Q_{n,1}$. Besides, taking limits in (4) and using again the size of derivatives of Laguerre polynomials, we get (b) for the polynomials $Q_{n,1}$.

If there are no more positive masses, since $Q_{n,1} = Q_n$ we have concluded the proof. Otherwise, suppose that the results (a) and (b) hold for the sequence of polynomials $\{Q_{n,s-1}\}_{n \geq 0}$ orthogonal with respect to the inner product

$$(p, q)_{s-1} = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty p(x)q(x)x^\alpha e^{-x} dx + M_{j_1} p^{(j_1)}(0)q^{(j_1)}(0) + \dots + M_{j_{s-1}} p^{(j_{s-1})}(0)q^{(j_{s-1})}(0),$$

where $j_1 < j_2 < \dots < j_{s-1}$ and all these masses are positive. Now, we have to prove the result for the polynomials $Q_{n,s}$ orthogonal with respect to

$$(p, q)_s = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty p(x)q(x)x^\alpha e^{-x} dx + M_{j_1} p^{(j_1)}(0)q^{(j_1)}(0) + \dots + M_{j_s} p^{(j_s)}(0)q^{(j_s)}(0),$$

where $M_{j_s} > 0$. Since $(p, q)_s = (p, q)_{s-1} + M_{j_s} p^{(j_s)}(0)q^{(j_s)}(0)$ we can work as before. Then the Fourier expansion of the polynomial $Q_{n,s}$ in the orthogonal basis $\{Q_{n,s-1}\}_{n \geq 0}$ leads to

$$Q_{n,s}(x) = Q_{n,s-1}(x) - M_{j_s} Q_{n,s}^{(j_s)}(0) K_{n-1,s-1}^{(0,j_s)}(x, 0),$$

where $K_{n,s-1}$ denotes the corresponding n th kernel for the sequence $\{Q_{n,s-1}\}$ and

$$K_{n,s-1}^{(k,h)}(x, y) = \sum_{i=0}^n \frac{Q_{i,s-1}^{(k)}(x) Q_{i,s-1}^{(h)}(y)}{(Q_{i,s-1}, Q_{i,s-1})_{s-1}}, \quad k, h \in \mathbb{N} \cup \{0\}.$$

Therefore, in the same way as in (3) and (4), we get

$$Q_{n,s}(x) = Q_{n,s-1}(x) - \frac{M_{j_s} Q_{n,s-1}^{(j_s)}(0)}{1 + M_{j_s} K_{n-1,s-1}^{(j_s,j_s)}(0, 0)} K_{n-1,s-1}^{(0,j_s)}(x, 0), \tag{8}$$

and

$$(Q_{n,s}, Q_{n,s})_s = (Q_{n,s-1}, Q_{n,s-1})_{s-1} + M_{j_s} \frac{(Q_{n,s-1}^{(j_s)}(0))^2}{1 + M_{j_s} K_{n-1,s-1}^{(j_s,j_s)}(0, 0)}. \tag{9}$$

Taking derivatives k times in (8) and evaluating at $x = 0$, we obtain

$$\frac{Q_{n,s}^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} = \frac{Q_{n,s-1}^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} \left[1 - \frac{M_{j_s} K_{n-1,s-1}^{(k,j_s)}(0, 0)}{1 + M_{j_s} K_{n-1,s-1}^{(j_s,j_s)}(0, 0)} \frac{Q_{n,s-1}^{(j_s)}(0)}{Q_{n,s-1}^{(k)}(0)} \right]. \tag{10}$$

Applying the Stolz criterion and the hypotheses (a) and (b) for $\{Q_{n,s-1}\}_{n \geq 0}$, we can obtain

$$K_{n-1,s-1}^{(k,j_s)}(0, 0) \cong \begin{cases} C_k n^{\alpha+k+j_s+1}, & \text{if } k \neq j_1, \dots, j_{s-1}; \\ C_k n^{j_s-k}, & \text{if } k = j_1, \dots, j_{s-1}, \end{cases} \tag{11}$$

where C_k is a nonzero constant. Indeed, for $k \neq j_1, \dots, j_{s-1}$,

$$\begin{aligned} \lim_n \frac{K_{n-1,s-1}^{(k,j_s)}(0, 0)}{n^{\alpha+k+j_s+1}} &= \lim_n \frac{Q_{n-1,s-1}^{(k)}(0) Q_{n-1,s-1}^{(j_s)}(0)}{(Q_{n-1,s-1}, Q_{n-1,s-1})_{s-1} (\alpha + k + j_s + 1) n^{\alpha+k+j_s}} \\ &= \lim_n \frac{Q_{n-1,s-1}^{(k)}(0)}{(L_{n-1}^\alpha)^{(k)}(0)} \lim_n \frac{Q_{n-1,s-1}^{(j_s)}(0)}{(L_{n-1}^\alpha)^{(j_s)}(0)} \lim_n \frac{(L_{n-1}^\alpha)^{(k)}(0) (L_{n-1}^\alpha)^{(j_s)}(0)}{\|L_{n-1}^\alpha\|^2 (\alpha + k + j_s + 1) n^{\alpha+k+j_s}}, \end{aligned} \tag{12}$$

and, for $k = j_1, \dots, j_{s-1}$,

$$\begin{aligned} \lim_n \frac{K_{n-1,s-1}^{(k,j_s)}(0, 0)}{n^{j_s-k}} &= \lim_n \frac{Q_{n-1,s-1}^{(k)}(0) Q_{n-1,s-1}^{(j_s)}(0)}{(Q_{n-1,s-1}, Q_{n-1,s-1})_{s-1} (j_s - k) n^{j_s-k-1}} \\ &= \lim_n \frac{(L_{n-1}^\alpha)^{(k)}(0) (L_{n-1}^\alpha)^{(j_s)}(0)}{\|L_{n-1}^\alpha\|^2 (j_s - k) n^{\alpha+k+j_s}} \lim_n n^{\alpha+2k+1} \frac{Q_{n-1,s-1}^{(k)}(0)}{(L_{n-1}^\alpha)^{(k)}(0)} \lim_n \frac{Q_{n-1,s-1}^{(j_s)}(0)}{(L_{n-1}^\alpha)^{(j_s)}(0)}. \end{aligned} \tag{13}$$

Then, from (10), (11) and the hypothesis for $Q_{n,s-1}$, we have

$$\frac{Q_{n,s}^{(j_s)}(0)}{(L_n^\alpha)^{(j_s)}(0)} = \frac{Q_{n,s-1}^{(j_s)}(0)}{(L_n^\alpha)^{(j_s)}(0)} \frac{1}{1 + M_{j_s} K_{n-1,s-1}^{(j_s,j_s)}(0,0)} \cong \frac{C_{j_s}}{n^{\alpha+2j_s+1}},$$

with C_{j_s} a nonzero constant. Moreover, for $k \neq j_s$, taking into account (12), (13) and the hypothesis for $Q_{n,s-1}$, we can deduce

$$\begin{aligned} \frac{K_{n-1,s-1}^{(k,j_s)}(0,0) Q_{n,s-1}^{(j_s)}(0)}{K_{n-1,s-1}^{(j_s,j_s)}(0,0) Q_{n,s-1}^{(k)}(0)} &= \frac{K_{n-1,s-1}^{(k,j_s)}(0,0) Q_{n,s-1}^{(j_s)}(0)}{K_{n-1,s-1}^{(j_s,j_s)}(0,0) (L_n^\alpha)^{(j_s)}(0)} \frac{(L_n^\alpha)^{(k)}(0)}{Q_{n,s-1}^{(k)}(0)} \frac{(L_n^\alpha)^{(j_s)}(0)}{(L_n^\alpha)^{(k)}(0)} \\ &\cong \frac{(L_n^\alpha)^{(j_s)}(0)}{(L_{n-1}^\alpha)^{(j_s)}(0)} \frac{(L_{n-1}^\alpha)^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} \begin{cases} \frac{\alpha+2j_s+1}{\alpha+k+j_s+1}, & \text{if } k \neq j_1, \dots, j_{s-1}; \\ \frac{\alpha+2j_s+1}{j_s-k}, & \text{if } k = j_1, \dots, j_{s-1}, \end{cases} \\ &\cong \begin{cases} \frac{\alpha+2j_s+1}{\alpha+k+j_s+1}, & \text{if } k \neq j_1, \dots, j_{s-1}; \\ \frac{\alpha+2j_s+1}{j_s-k}, & \text{if } k = j_1, \dots, j_{s-1}. \end{cases} \end{aligned}$$

Thus, taking limits in (10) and (9), we get (a) and (b) for the polynomials $Q_{n,s}$, i.e.

$$\frac{Q_{n,s}^{(k)}(0)}{(L_n^\alpha)^{(k)}(0)} \cong \begin{cases} \frac{C_k}{n^{\alpha+2k+1}}, & \text{if } k = j_1, \dots, j_s; \\ C_k, & \text{otherwise,} \end{cases}$$

and

$$(Q_{n,s}, Q_{n,s})_s \cong \|L_n^\alpha\|^2.$$

Hence the result follows. \square

Observe that the part (a) of Lemma 1 is also true for the ratio of the corresponding orthonormal polynomials, and therefore there exists

$$\lim_n \frac{q_n^{(k)}(0)}{(l_n^\alpha)^{(k)}(0)} = \begin{cases} 0, & \text{for } k \text{ such that } M_k > 0; \\ C_k \neq 0, & \text{otherwise.} \end{cases} \tag{14}$$

Consider the following representation of the orthonormal polynomials q_n in terms of the orthonormal Laguerre polynomials l_n^α (see [8, Section 9])

$$q_n(x) = \sum_{j=0}^{N+1} b_j(n) x^j l_{n-j}^{\alpha+2j}(x). \tag{15}$$

For the inner product (1) with $N = 1$, the coefficients $b_j(n)$ was explicitly obtained in [9], and their estimation was essential to obtain the result in [4].

Now in the general case, using Lemma 1, we can prove that there is always limit of the connection coefficients $b_j(n)$ for an arbitrary N .

Lemma 2. Let $\{b_j(n)\}_0^{N+1}$ be the coefficients in formula (15). Then, there exists

$$\lim_n b_j(n) = b_j \in \mathbb{R}, \quad j \in \{0, \dots, N + 1\}.$$

Moreover, the first index j such that $b_j \neq 0$ corresponds with the first j such that $M_j = 0$ in the inner product (1). (We understand that if all the masses are positive, then the unique coefficient b_j different from zero is the last one.)

Proof. Taking derivatives k times in (15) and evaluating at $x = 0$, we deduce

$$\frac{q_n^{(k)}(0)}{(l_n^\alpha)^{(k)}(0)} = \sum_{j=0}^k b_j(n) \binom{k}{j} j! A_j(k, n), \quad k \in \{0, \dots, N + 1\}, \tag{16}$$

where $A_0(k, n) = 1$ and

$$A_j(k, n) = \frac{(l_{n-j}^{\alpha+2j})^{(k-j)}(0)}{(l_n^\alpha)^{(k)}(0)} \cong \frac{(-1)^j \Gamma(\alpha + k + 1)}{\Gamma(\alpha + k + j + 1)} \left(\frac{\Gamma(\alpha + 2j + 1)}{\Gamma(\alpha + 1)} \right)^{1/2}. \tag{17}$$

Since there exists $\lim_n A_j(k, n) \neq 0$, applying recursively (14) and (16) we can assure there exists $\lim_n b_j(n) = b_j$, $j \in \{0, \dots, N + 1\}$. More precisely, for $k = 0$ we have

$$\lim_n b_0(n) = \lim_n \frac{q_n(0)}{l_n^\alpha(0)} = b_0 = \begin{cases} 0, & \text{if } M_0 > 0; \\ C \neq 0, & \text{if } M_0 = 0. \end{cases}$$

Now, from (16) for $k = 1$, (14) and (17) we get

$$\lim_n b_1(n) = \lim_n \frac{1}{A_1(1, n)} \left(\frac{q_n'(0)}{(l_n^\alpha)'(0)} - b_0(n) \right) = b_1.$$

Observe that

$$b_1 = \begin{cases} 0, & \text{if } M_0 > 0 \text{ and } M_1 > 0; \\ C \neq 0, & \text{if } M_0 > 0 \text{ and } M_1 = 0. \end{cases}$$

In this way, recursively, if $M_0 M_1 \dots M_i > 0$ and $M_{i+1} = 0$ we can assure that

$$b_j = \begin{cases} 0, & \text{if } 0 \leq j \leq i; \\ C \neq 0, & \text{if } j = i + 1, \end{cases}$$

and we obtain the result. \square

As a consequence of the above lemma, we can establish a Mehler–Heine type formula for general discrete Laguerre–Sobolev orthonormal polynomials. This formula shows how the presence of the masses in the discrete part of the inner product changes the asymptotic behavior around the origin. Moreover, it supplies information on the location and asymptotic distribution of the zeros of the polynomials in terms of the zeros of known special functions.

We recall the corresponding formula for orthonormal Laguerre polynomials (see [14])

$$\lim_n \frac{l_n^\alpha(x/(n+k))}{n^{\alpha/2}} = \sqrt{\Gamma(\alpha+1)} x^{-\alpha/2} J_\alpha(2\sqrt{x}) \tag{18}$$

uniformly on compact subsets of \mathbb{C} and uniformly for $k \in \mathbb{N} \cup \{0\}$, where J_α is the Bessel function of the first kind.

Proposition 1. *The polynomials q_n satisfy the following Mehler–Heine type formula:*

$$\lim_n \frac{q_n(x/n)}{n^{\alpha/2}} = \sqrt{\Gamma(\alpha+1)} \sum_{j=0}^{N+1} b_j x^{-\alpha/2} J_{\alpha+2j}(2\sqrt{x}) \tag{19}$$

uniformly on compact subsets of \mathbb{C} .

Proof. The proof is a straightforward consequence of formula (15), Lemma 2 and (18). \square

Remark. According to Lemma 2, the first Bessel function which appears in (19) corresponds with the first index j such that $M_j = 0$, in the inner product (1). We want to highlight that this result generalizes the one obtained in [1, Theorem 3], where the authors only deal with inner products with a unique “gap” in the discrete part.

The above proposition allows us to deduce a lower estimate of $\|q_n\|_{L_p(x^\beta dx)}$, for $\beta = \alpha$ and $\beta = \alpha p/2$, that will play an important role in the proof of Theorem 1.

Proposition 2. *Let $1 \leq p \leq \infty$. Then, the following statements hold:*

For $\alpha > -1/2$

$$\|q_n\|_{L_p(x^\alpha dx)} \geq C \begin{cases} n^{-1/4} (\log(n+1))^{1/p}, & \text{if } p = \frac{4\alpha+4}{2\alpha+1}; \\ n^{\alpha/2 - (\alpha+1)/p}, & \text{if } \frac{4\alpha+4}{2\alpha+1} < p \leq \infty, \end{cases}$$

and for $\alpha > -2/p$ if $1 \leq p < \infty$ and $\alpha \geq 0$ if $p = \infty$

$$\|q_n\|_{L_p(x^{\alpha p/2} dx)} \geq C \begin{cases} n^{-1/4} (\log(n+1))^{1/p}, & \text{if } p = 4; \\ n^{-1/p}, & \text{if } 4 < p \leq \infty, \end{cases}$$

where C is an absolute positive constant.

Proof. Assume $1 \leq p < \infty$. Then,

$$\|q_n\|_{L_p(x^\beta dx)}^p = \int_0^\infty |q_n(x)e^{-x/2}|^p x^\beta dx > \int_0^{1/\sqrt{n}} |q_n(x)e^{-x/2}|^p x^\beta dx \geq Cn^{-\beta-1} \int_0^{\sqrt{n}} |q_n(t/n)|^p t^\beta dt.$$

According to formula (19), $\exists n_0 \in \mathbb{N}$ such that $\forall n \geq n_0$

$$\int_0^{\sqrt{n}} |q_n(t/n)|^p t^\beta dt \geq Cn^{p\alpha/2} \int_0^{\sqrt{n}} \left| \sum_{j=0}^{N+1} b_j t^{-\alpha/2} J_{\alpha+2j}(2\sqrt{t}) \right|^p t^\beta dt$$

and therefore $\forall n \geq n_0$

$$\|q_n\|_{L_p(x^\beta dx)}^p \geq Cn^{p\alpha/2-\beta-1} \int_0^{2n^{1/4}} u^{2\beta-p\alpha+1} \left| \sum_{j=0}^{N+1} b_j J_{\alpha+2j}(u) \right|^p du.$$

Working as Stempak in [13, Lemma 2.1], we can prove that for $\alpha > -1$, and $\lambda > -1 - \alpha p$

$$\int_0^{2n^{1/4}} u^\lambda \left| \sum_{j=0}^{N+1} b_j J_{\alpha+2j}(u) \right|^p du \sim \begin{cases} 1, & \text{if } \lambda < p/2 - 1; \\ \log(n+1), & \text{if } \lambda = p/2 - 1. \end{cases}$$

Thus, if $1 \leq p < \infty$, we obtain the first and the second result for $\beta = \alpha$ and $\beta = p\alpha/2$ respectively. The results for $p = \infty$ can be deduced from the previous one by passing to the limit when p goes to ∞ . \square

It is worth noticing that these lower bounds are sharp in the following sense.

Proposition 3. Let $1 \leq p \leq \infty$. Then:

For $\alpha \geq 0$,

$$\|q_n\|_{L_p(x^\alpha dx)} \sim \begin{cases} n^{-1/4}(\log(n+1))^{1/p}, & \text{if } p = \frac{4\alpha+4}{2\alpha+1}; \\ n^{\alpha/2-(\alpha+1)/p}, & \text{if } \frac{4\alpha+4}{2\alpha+1} < p \leq \infty, \end{cases}$$

and for $\alpha > -2/p$ if $1 \leq p < \infty$ and $\alpha \geq 0$ if $p = \infty$,

$$\|q_n\|_{L_p(x^{\alpha p/2} dx)} \sim \begin{cases} n^{-1/4}(\log(n+1))^{1/p}, & \text{if } p = 4; \\ n^{-1/p}, & \text{if } 4 < p \leq \infty. \end{cases}$$

Proof. From Lemma 1 of [11] it can be deduced that for $\alpha \geq 0$

$$\int_0^\infty |x^j l_n^{\alpha+2j}(x)e^{-x/2}|^p x^\alpha dx \sim \begin{cases} n^{-p/4} \log(n+1), & \text{if } p = \frac{4\alpha+4}{2\alpha+1}; \\ n^{\alpha p/2-(\alpha+1)}, & \text{if } \frac{4\alpha+4}{2\alpha+1} < p \leq \infty, \end{cases}$$

and for $\alpha > -2/p$ if $1 \leq p < \infty$ and $\alpha \geq 0$ if $p = \infty$

$$\int_0^\infty |x^j l_n^{\alpha+2j}(x)e^{-x/2} x^{\alpha/2}|^p dx \sim \begin{cases} n^{-p/4} \log(n+1), & \text{if } p = 4; \\ n^{-1}, & \text{if } 4 < p \leq \infty. \end{cases}$$

Thus, using the representation formula for the polynomials q_n (see (15)), and the fact that the connection coefficients are bounded (see Lemma 2), we get one of the two inequalities. The other one has been proved in Proposition 2 and therefore the result follows. \square

3. A Cohen type inequality

In this section we prove a Cohen type inequality for the Fourier expansions in terms of discrete Laguerre–Sobolev orthonormal polynomials with an arbitrary (finite) number of mass points. So we extend the result due to Fejzullahu and Marcellán which deals with a discrete Laguerre–Sobolev inner product with at most two masses in the discrete part (see [4]).

Proof of Theorem 1. Let us consider the following test functions which were already used in [12] and later in [4]

$$g_n^{\alpha,j}(x) = x^j \left[L_n^{\alpha+j}(x) - \sqrt{\frac{(n+1)(n+2)}{(n+\alpha+j+1)(n+\alpha+j+2)}} L_{n+2}^{\alpha+j}(x) \right],$$

with $j \in \mathbb{N} \setminus \{1, \dots, N\}$. Notice that

$$(g_n^{\alpha,j})^{(i)}(0) = 0, \quad i = 0, \dots, N. \tag{20}$$

These functions can be written as (see formula (2.15) in [12])

$$g_n^{\alpha,j}(x) = \sum_{m=0}^{j+2} a_{m,j}(\alpha, n) L_{n+m}^{\alpha}(x) \tag{21}$$

with

$$a_{0,j}(\alpha, n) = \frac{\Gamma(n + \alpha + j + 1)}{\Gamma(n + \alpha + 1)} \cong n^j.$$

From (20), (21), and $0 \leq k \leq n$, we have

$$\begin{aligned} \widehat{g_n^{\alpha,j}}(k) &= \langle g_n^{\alpha,j}, q_k \rangle_S = \frac{1}{\Gamma(\alpha + 1)} \int_0^\infty g_n^{\alpha,j}(x) q_k(x) e^{-x} x^\alpha dx \\ &= \frac{1}{\Gamma(\alpha + 1)} \sum_{m=0}^{j+2} a_{m,j}(\alpha, n) \int_0^\infty L_{n+m}^{\alpha}(x) q_k(x) e^{-x} x^\alpha dx. \end{aligned}$$

By the orthogonality of Laguerre polynomials, we obtain

$$\widehat{g_n^{\alpha,j}}(k) = \begin{cases} 0, & \text{if } 0 \leq k \leq n - 1; \\ \frac{1}{\Gamma(\alpha+1)} a_{0,j}(\alpha, n) \int_0^\infty L_n^{\alpha}(x) q_n(x) e^{-x} x^\alpha dx, & \text{if } k = n. \end{cases}$$

Thus, from Lemma 1(b), the estimate of $a_{0,j}(\alpha, n)$ and the value of the norm of Laguerre polynomials (see (2)), we can deduce

$$\begin{aligned} \widehat{g_n^{\alpha,j}}(n) &= \frac{1}{\Gamma(\alpha + 1)} a_{0,j}(\alpha, n) \int_0^\infty L_n^{\alpha}(x) \frac{Q_n(x)}{\langle Q_n, Q_n \rangle_S^{1/2}} e^{-x} x^\alpha dx \\ &= a_{0,j}(\alpha, n) \frac{\|L_n^{\alpha}\|^2}{\langle Q_n, Q_n \rangle_S^{1/2}} \cong a_{0,j}(\alpha, n) \|L_n^{\alpha}\| \cong \frac{n^{j+\alpha/2}}{\sqrt{\Gamma(\alpha + 1)}}. \end{aligned}$$

Observe that Q_n and L_n^{α} have always equivalent norms, and, therefore this estimation does not depend neither on the number of positive masses, nor on the existence or non-existence of any gap in the inner product.

Applying the operator $T_n^{\alpha,S}$ to the functions $g_n^{\alpha,j}$, we get

$$T_n^{\alpha,S}(g_n^{\alpha,j}) = c_{n,n} \widehat{g_n^{\alpha,j}}(n) q_n,$$

and therefore

$$\begin{aligned} \|T_n^{\alpha,S}\|_{[S_p^\beta]} &\geq (\|g_n^{\alpha,j}\|_{S_p^\beta})^{-1} \|T_n^{\alpha,S}(g_n^{\alpha,j})\|_{S_p^\beta} = (\|g_n^{\alpha,j}\|_{S_p^\beta})^{-1} |c_{n,n}| |\widehat{g_n^{\alpha,j}}(n)| \|q_n\|_{S_p^\beta} \\ &\geq (\|g_n^{\alpha,j}\|_{S_p^\beta})^{-1} |c_{n,n}| |\widehat{g_n^{\alpha,j}}(n)| \|q_n\|_{L_p(x^\beta dx)}. \end{aligned}$$

On the other hand, for $j > \alpha - 1/2 - 2(\alpha + 1)/p$ we have

$$\|g_n^{\alpha,j}\|_{S_p^\beta} \leq c \begin{cases} n^{j-1/2+(\alpha+1)/p}, & \text{if } \beta = \alpha; \\ n^{\alpha/2+j-1/2+1/p}, & \text{if } \beta = p\alpha/2 \end{cases}$$

(see formula (3.3) and formula (1.19), (2.12) in [12] respectively). Thus, by Proposition 2 we get:

For $\beta = \alpha$ with $\alpha > -1/2$

$$\|T_n^{\alpha,S}\|_{[S_p^{\alpha}]} \geq C|c_{n,n}| \begin{cases} (\log(n+1))^{\frac{2\alpha+1}{4\alpha+4}}, & \text{if } p = q_0; \\ n^{\alpha+1/2-2(\alpha+1)/p}, & \text{if } q_0 < p \leq \infty. \end{cases}$$

For $\beta = p\alpha/2$ with $\alpha > -2/p$ if $1 \leq p < \infty$ and $\alpha \geq 0$ if $p = \infty$,

$$\|T_n^{\alpha,S}\|_{[S_p^{p\alpha/2}]} \geq C|c_{n,n}| \begin{cases} (\log(n+1))^{1/4}, & \text{if } p = 4; \\ n^{1/2-2/p}, & \text{if } 4 < p \leq \infty. \end{cases}$$

Hence, by duality the theorem follows. \square

Remark. In particular, for $M_i = 0$, $i = 0, \dots, N$, the above theorem extends Theorem 1 in [12] to negative values of α .

In the particular case of $c_{k,n} = 1$, $k = 0, \dots, n$, the operator $T_n^{\alpha,S}$ is the n th partial sum of the Fourier expansion, so, we can assure the following result.

Corollary 1. If p is outside the Pollard interval (p_0, q_0) , we have

$$\|S_n\|_{[S_p^{\beta}]} \rightarrow \infty, \quad n \rightarrow \infty$$

where S_n denotes the n th partial sum of the Fourier expansion.

References

- [1] M. Alfaro, J.J. Moreno-Balcázar, A. Peña, M.L. Rezola, A new approach to the asymptotics of Sobolev type orthogonal polynomials, J. Approx. Theory 163 (2011) 460–480.
- [2] R. Askey, S. Wainger, Mean convergence of expansions in Laguerre and Hermite series, Amer. J. Math. 87 (1965) 695–708.
- [3] P.J. Cohen, On a conjecture of Littlewood and idempotent measures, Amer. J. Math. 82 (1960) 191–212.
- [4] B.Xh. Fejzullahu, F. Marcellán, A Cohen type inequality for Laguerre–Sobolev expansions, J. Math. Anal. Appl. 352 (2009) 880–889.
- [5] S. Giulini, P.M. Sordi, G. Travaglini, A Cohen type inequality for compact Lie groups, Proc. Amer. Math. Soc. 77 (1979) 359–364.
- [6] G.H. Hardy, J.E. Littlewood, A new proof of a theorem on rearrangements, J. Lond. Math. Soc. 23 (1948) 163–168.
- [7] G. Klambauer, Aspects of Calculus, Springer-Verlag, New York, 1986.
- [8] R. Koekoek, Generalizations of Laguerre polynomials, J. Math. Anal. Appl. 153 (1990) 576–590.
- [9] R. Koekoek, H.G. Meijer, A generalization of Laguerre polynomials, SIAM J. Math. Anal. 24 (1993) 768–782.
- [10] B. Muckenhoupt, Mean convergence of Hermite and Laguerre series II, Trans. Amer. Math. Soc. 147 (1970) 433–460.
- [11] C. Markett, Mean Cesàro summability of Laguerre expansions and norm estimates with shifted parameter, Anal. Math. 8 (1982) 19–37.
- [12] C. Markett, Cohen type inequalities for Jacobi, Laguerre and Hermite expansions, SIAM J. Math. Anal. 14 (1983) 819–833.
- [13] K. Stempak, On convergence and divergence of Fourier–Bessel series, Electron. Trans. Numer. Anal. 14 (2002) 223–235.
- [14] G. Szegő, Orthogonal Polynomials, fourth edition, Amer. Math. Soc. Colloq. Publ., vol. 23, Amer. Math. Soc., Providence, RI, 1975.